AQA

Surname	
Other Names	
Centre Number	
Candidate Number	
Candidate Signature	

A-level BIOLOGY

Paper 2

7402/2

Monday 11 June 2018 Afternoon

Time allowed: 2 hours

For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

At the top of the page, write your surname and other names, your centre number, your candidate number and add your signature.



www.xtrapapers.com

BLANK PAGE



INSTRUCTIONS

- Use black ink or black ball-point pen.
- Answer ALL questions.
- You must answer the questions in the space provided. Do not write on blank pages.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

INFORMATION


- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 91.

DO NOT TURN OVER UNTIL TOLD TO DO SO



Answer ALL questions in the spaces provided.

0 1 Heat stress is a condition that often occurs in plants exposed to high temperatures for a prolonged period of time. Heat stress is a major factor in limiting the rate of photosynthesis.



1 Heat stress decreases the light-dependent reaction of photosynthesis.

Explain why this leads to a decrease in the LIGHT-INDEPENDENT REACTION. [2 marks]



01.2	Another effect of heat stress is a decrease in the activity of the enzyme rubisco. A decrease in the activity of an enzyme means that the rate of the reaction it catalyses becomes slower.
	A decrease in the activity of the enzyme rubisco would limit the rate of photosynthesis.
	Explain why. [2 marks]
01.3	Where precisely is rubisco found in a cell? [1 mark]
[Turn ove	r]



Scientists investigated the effect of temperature on the activity of two enzymes isolated from the leaf cells of cotton plants.

- Rubisco
- Rubisco activase an enzyme that activates rubisco

FIGURE 1 and FIGURE 2 show their results.







FIGURE 2 Rubisco activase activity / arbitrary units 0.5 0.4



01.4 The scientists concluded that heat stress reduces the activity of rubisco in plant leaves by affecting rubisco activase.

Use all the information to evaluate their conclusion. [4 marks]



www.xtrapapers.com

BLANK PAGE



[Tur	'n o	ver]



02.1 There are different types of gene mutation.

Put a tick (\checkmark) in the box next to the statement which describes INCORRECTLY the effect of the mutation in an exon of a gene. [1 mark]



A substitution may not result in a change to the encoded amino acid.



An inversion will result in a change in the number of DNA bases.



A deletion will result in a frame shift.



An addition will result in a frame shift.

02.2 Describe how alterations to tumour suppressor genes can lead to the development of tumours. [3 marks]





BLANK PAGE

02.3 A type of malignant tumour cell divides every 8 hours.

Starting with one of these cells, how many tumour cells will be present after 4 weeks? Assume none of these cells will die.

Give your answer in standard form. [2 marks]

Answer =




Yeast cells can respire aerobically or anaerobically. A student used the apparatus shown in FIGURE 3 to measure the rate of respiration in yeast.

She:

- positioned the flask in a water bath so that the yeast culture reached a constant temperature
- then left the apparatus for one hour before starting her investigation.

FIGURE 3





03.1 Suggest ONE reason why it was important that the student left the apparatus for one hour after the yeast culture reached a constant temperature. [1 mark]



Explain why it moved to the right. [2 marks]



BLANK PAGE

_			
_			
_			
_			
_			
[Turn over]		



03.3 The student found that the coloured liquid moved 1.5 cm in 24 hours. The diameter of the lumen (hole) of the capillary tubing was 1 mm.

The volume of a capillary tubing is given by $\pi r^2 l$, where π is 3.14 and l = length.


Calculate the volume of gas produced in cm³ hour⁻¹. Show your working. [2 marks]

Answer =

cm³ hour−1








03.4 Explain why a log scale is used to record the number of cells. [1 mark]



03.5 Many yeast cells die during the death phase.

Suggest ONE reason why. [1 mark]



 $X_t = X_0 e^{rt}$

 X_t = the population after a certain time X_0 = the population at the start e = 2.72 (base of natural logarithm) r = growth rate t = time period in hours over which r applies

A population of 2000 yeast cells was left for 10 hours.

The value for the growth rate was 0.5



Assuming no yeast cells died, calculate the predicted size of the population after 10 hours. Show your working. [2 marks]


Answer =

9



Scientists investigated the process of succession on sand dunes. They measured the percentage cover of different species of plants on sand dunes of different ages. Some of the results the scientists obtained are shown in FIGURE 5.

FIGURE 5





04.1 Describe how you would determine the mean percentage cover for beach grass on a sand dune. [3 marks]




BLANK PAGE

The scientists concluded that the results
shown in FIGURE 5 were due to succession
taking place.

Use FIGURE 5 to explain why the scientists reached this conclusion. [4 marks]








changed during succession. Some of the results the scientists obtained are shown in FIGURE 6.



FIGURE 6





BLANK PAGE

04.4 Using evidence from FIGURE 6, what can you conclude about the net primary productivity (NPP) in the sand dunes that are older than 1000 years?

Explain your answer. [2 marks]

[Turn over]



Furosemide and CVT are drugs used to remove excess fluid from the body. Scientists investigated the effect of these drugs on the volume of urine produced by human volunteers. The scientists used the following method.

- They divided volunteers into three groups, A, B and C at random.
- They gave all the volunteers the same food for 3 days.
- After 3 days, they gave the volunteers in group A a tablet containing furosemide, the volunteers in group B a tablet containing CVT and the volunteers in group C a placebo (a tablet that did NOT contain either drug).
- They then found the mean volume of urine produced by each group in the 4 hours after taking the tablets.

Some of the results the scientists obtained are shown in TABLE 1.

A value of \pm 2 standard deviations from the mean includes over 95% of the data.



TABLE 1

Group	Mean volume of urine produced in 4 hours / cm ³ (\pm 2 standard deviations)
A	1980
(furosemide)	(± 152)
B	1201
(CVT)	(± 119)
C	312
(placebo)	(± 57)

05.1 All the volunteers were given the same food for 3 days.

Suggest and explain ONE reason why they were given the same food. [2 marks]









05.2 Using TABLE 1, what can you conclude about the effectiveness of furosemide and CVT in the removal of excess fluid from the body? [2 marks]



05.3 Furosemide is sometimes used to treat high blood pressure.

Suggest how furosemide would cause a decrease in blood pressure. [1 mark]



05.4 Furosemide inhibits the absorption of sodium and chloride ions from the filtrate produced in the nephrons.

Explain how furosemide causes an increase in the volume of urine produced. [3 marks]



The scientists also measured the mean rate of flow of blood plasma into the kidneys.

The results the scientists obtained are shown in TABLE 2.

TABLE 2

Group	Mean rate of flow of blood plasma into the kidneys / cm ³ min ⁻¹
A (furosemide)	380
B (CVT)	342
C (placebo)	295



05.5 The mean rate of flow of blood plasma is 60% of the mean rate of blood flow into the kidneys.

How much greater is the flow of blood into the kidneys with furosemide than with group C (placebo) over the 4 hours of the investigation? Give your answer in cm³. [1 mark]

Answer =

cm³



06.1 In genetic crosses, the observed phenotypic ratios obtained in the offspring are often NOT the same as the expected ratios.

Suggest TWO reasons why. [2 marks]



In tomato plants, the genes for height and for the type of leaf are on the same homologous pair of chromosomes. The allele T, for a tall plant, is dominant to the allele t, for a dwarf plant. The allele M, for normal leaves, is dominant to the allele m, for mottled leaves.

A biologist carried out crosses between parent plants heterozygous for both genes and examined the offspring produced. The position of the two alleles for both genes was the same in each parent plant as shown in FIGURE 7 below. The phenotypes and number of offspring produced are shown in TABLE 3 on page 40.

FIGURE 7







TABLE 3

Phenotype of offspring	Number of offspring
Tall plants and normal leaves	1860
Tall plants and mottled leaves	68
Dwarf plants and normal leaves	57
Dwarf plants and mottled leaves	580

06.2 What would be the genotype of the offspring with dwarf plants and mottled leaves? [1 mark]



06.3 Use the information provided to explain the results in TABLE 3. [3 marks]



41



BLANK PAGE

4 2

06.4 Complete TABLE 4 to show the expected ratio of phenotypes if the same cross had been carried out but the genes for height of plant and for the type of leaf were on different homologous pairs of chromosomes. [2 marks]

TABLE 4

Phenotype of offspring	Ratio of offspring

[Turn over]

8




07.1 Dopamine is a neurotransmitter released in some synapses in the brain. The transmission of dopamine is similar to that of acetylcholine.

Dopamine stimulates the production of nerve impulses in postsynaptic neurones.


Describe how.


Do NOT include in your answer the events leading to the release of dopamine and the events following production of nerve impulses at postsynaptic neurones. [3 marks]














Chromatin immunoprecipitation is one method to determine where a transcription factor binds to DNA. The principle behind this procedure is shown in FIGURE 8.

FIGURE 8






08.2 Explain why the antibody binds to the transcription factor. [2 marks]

08.3	Use FIGURE 8 to explain what 'precipitated DNA' consists of. [1 mark]



Soybeans are used in a number of processed foods. However, soybeans contain a protein known as P34 that causes an allergic response in some people. Scientists have created transgenic soybeans that produce singlestranded cDNA, which prevents transcription of the P34 gene. They used recombinant plasmids as vectors to transform soybean cells. After they had screened these cells for production of the P34 protein, they cultured the transformed cells to form soybean plants.

08.4 Suggest how single-stranded cDNA could prevent transcription of the P34 gene. [1 mark]



0 8 . 5 Describe the roles of TWO named types of enzymes used to insert DNA fragments into plasmids. [2 marks]



	Type of enzyme
	Role
	Type of enzyme
	Role
[Turn ove	r]



BLANK PAGE

5 2

0 8 . 6	The soybean cells were screened for the
	presence of the P34 protein. This process involved the use of gel electrophoresis to separate proteins extracted from soybean cells.

Suggest TWO features of the structure of different proteins that enable them to be separated by gel electrophoresis. [2 marks]

1			
2			







Lactose is the main sugar in milk and is hydrolysed by the enzyme lactase. Lactase is essential to newborn mammals as milk is their only source of food. Most mammals stop producing lactase when they start feeding on other food sources. Humans are an exception to this because some continue to produce lactase as adults. The ability to continue producing lactase is known as lactase persistence (LP) and is controlled by a dominant allele. A number of hypotheses based on different selection pressures have been put forward to explain LP in humans.

09.1 One hypothesis for LP in humans suggests that the selective pressure was related to some human populations farming cattle as a source of milk.

Describe how farming cattle as a source of milk could have led to an increase in LP. [4 marks]



55



09.2 Use the information provided to explain why the number of people showing LP would RAPIDLY increase once selection for this condition had been established. [2 marks]



09.3 Lactase persistence is caused by a mutation in DNA. This mutation does NOT occur in the gene coding for lactase.

Suggest and explain how this mutation causes LP. [2 marks]



8



10

Read the following passage.

Complete achromatopsia is a form of complete colour blindness. It is caused by having only rods and no functional cone cells. People with complete achromatopsia have difficulty in seeing 5 detail. Complete achromatopsia is caused by an autosomal recessive allele and is usually very rare in populations with only one in 40 000 being affected. However on the Pacific island of Pingelap ten percent 10 of the population are affected.

One form of red-green colour blindness is
caused by a sex-linked recessive allele
which affects more men than women.People with this red-green colour15blindness are unable to distinguish
between red and green, and also between
other colours. They have green-sensitive
cones but the photoreceptive pigment
they contain does not function.20

Scientists investigated the use of gene therapy to correct red-green colour blindness in monkeys. They injected viruses containing the gene for the greensensitive pigment directly into the eyes of 25 the monkeys. Although the monkeys maintained two years of colour vision, there is debate on whether this form of gene therapy is worthwhile. No clinical




trials of this procedure have been carried 30 out on humans. Current research into the treatment of red-green colour blindness involves the use of induced pluripotent stem cells (iPS cells). The use of iPS cells could have advantages over the use of 35 gene therapy.

Use the information in the passage and your own knowledge to answer the following questions.

10.1 People with complete achromatopsia have difficulty in seeing detail (lines 4-6).

Explain why. [3 marks]







Answer =	%

10.3 Red-green colour blindness affects more men than women (lines 12–14).

Explain why. [2 marks]



10.4 People with red-green colour blindness are unable to distinguish between red and green, and also between other colours (lines 15–18).

Explain why. [3 marks]				



10.5 Current research into the treatment of redgreen colour blindness involves the use of induced pluripotent stem cells (iPS cells) (lines 31–34).

Suggest how iPS cells could correct red-green colour blindness. [2 marks]



10.6 The use of iPS cells could have advantages over the use of gene therapy to correct red-green colour blindness (lines 34–36).

Using the information from the passage, suggest and explain reasons why. [3 marks]





END	OF	QUESTIONS	

15

6	5
---	---

66

There are no questions printed on this page

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
TOTAL		

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

IB/M/Jun18/IK/7402/2/E3

