

Please write clearly in	block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature			

AS BIOLOGY

Paper 1

Thursday 24 May 2018

Afternoon

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.


Information

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 75.

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
TOTAL		










Do not write outside the

box

0 1 . 3

Figure 2 shows a photograph of part of a mitochondrion from a mouse liver cell taken using a transmission electron microscope at \times 62 800 magnification.



Produce a scientific drawing of the mitochondrion in Figure 2 in the box below.

Label the following parts of the mitochondrion on your drawing.

- Matrix
- Crista

[4 marks]



Λ	
4	

02.1	What is a monomer? [1 mark]
02.2	Lactulose is a disaccharide formed from one molecule of galactose and one molecule of fructose.
	Other than both being disaccharides, give one similarity and one difference between the structures of lactulose and lactose.
	[2 marks]
	Similarity
	Difference



Do not write outside the box

0 2 . 3	Following digestion and absorption of food, the undigested remains are processed to	Do not write outside the box
	form faeces in the parts of the intestine below the ileum.	
	The faeces of people with constipation are dry and hard. Constipation can be treated by drinking lactulose. Lactulose is soluble, but is not digested or absorbed in the human intestine.	
	Use your knowledge of water potential to suggest why lactulose can be used to help people suffering from constipation.	
	[2 marks]	
02.4	Lactulose can also be used to treat people who have too high a concentration of hydrogen ions (H^{+}) in their blood.	
	The normal range for blood H * concentration is 3.55 \times 10 $^{-8}$ to 4.47 \times 10 $^{-8}$ mol dm $^{-3}$	
	A patient was found to have a blood $H^{\scriptscriptstyle +}$ concentration of 2.82 \times $10^{^{-7}}mol~dm^{^{-3}}$	
	Calculate the minimum percentage decrease required to bring the patient's blood H ⁺ concentration into the normal range.	
	[2 marks]	
		[]
	A	<u> </u>
1	Answer =	



Г


0 3 . 1	Draw and label a single DNA nucleotide.	Do not write outside the box
0 3.1	[2 marks]	
0 3.2	Cive two features of DNA and evolution how each and is important in the	
0 3.2	Give two features of DNA and explain how each one is important in the	
	semi-conservative replication of DNA. [2 marks]	
	semi-conservative replication of DNA.	
	semi-conservative replication of DNA. [2 marks]	
	semi-conservative replication of DNA. [2 marks]	
	semi-conservative replication of DNA. [2 marks]	
	semi-conservative replication of DNA. [2 marks]	
	semi-conservative replication of DNA. [2 marks] 1	
	semi-conservative replication of DNA. [2 marks]	
	semi-conservative replication of DNA. [2 marks] 1	
	semi-conservative replication of DNA. [2 marks] 1	
	semi-conservative replication of DNA. [2 marks] 1	
	semi-conservative replication of DNA. [2 marks] 1	
	semi-conservative replication of DNA. [2 marks] 1	
	semi-conservative replication of DNA. [2 marks] 1	



Do not write outside the 0 3 . 3 box Replication of mitochondrial DNA (mtDNA) is different from that of nuclear DNA. The replication of the second strand of mtDNA only starts after two-thirds of the first strand of mtDNA has been copied. A piece of mtDNA is 16 500 base pairs long and is replicated at a rate of 50 nucleotides per second. Tick (\checkmark) the box that shows how long it would take to copy this mtDNA. [1 mark] Α 330 seconds В 440 seconds С 550 seconds D 660 seconds 5 Turn over for the next question



Turn over ►





04.3	Name the process by which prokaryotic cells divide. [1 mark]	Do not write outside the box
04.4	Some prokaryotic cells can divide every 30 minutes. A liquid culture contained a starting population of 1.35×10^4 cells. Assuming each cell divides every 30 minutes, calculate how many cells there will be after 3 hours. Assume no cells die during this time. [2 marks]	
	Answer =	7
	Turn over for the next question	
	Turn over ►	



IB/M/Jun18/7401/1


0 5	Figure 4 represents a triglyceride.	Do not write outside the box
	Figure 4	
	Q	
	P Q	
	Q	
0 5.1	Name the molecules represented in Figure 4 by: [2 marks]	
	Box P	
	Box Q	
0 5.2	Name the type of bond between P and Q in Figure 4 . [1 mark]	



[2 marks]


Do not write outside the box

5




Turn over ►

IB/M/Jun18/7401/1










Turn over ►





Do not write outside the box

06.3	D. erecta and D. willistoni are closely related species but different species.
	Describe evidence from Figure 6 that supports this statement. [2 marks]
0 6.4	The scientists repeated their experiments, using female fruit flies as the visual stimulus. When a male and female <i>D. willistoni</i> were together, their songs led to mating.
	When two female <i>D.willistoni</i> were together, their songs did not lead to any attempt to mate.
	Use information from Figure 6 to suggest why the two females did not attempt to mate.
	[2 marks]

1 5


Turn over ►

0 7.1	What is the proteome of a cell? [1 mark]	Do not outside box
0 7.2	Give two structural differences between a molecule of messenger RNA (mRNA) and a molecule of transfer RNA (tRNA). [2 marks]	
	1	
	2	











0 8.1	State and explain the property of water that can help to buffer changes in temperature.	[2 marks]	Do not write outside the box
0 8.2	Water is used to hydrolyse ATP. Name the two products of ATP hydrolysis.	[1 mark]	
	12		



Do not write

box

Hydrolysis of ATP is catalysed by the enzyme ATP hydrolase.

A student investigated the effect of ATP concentration on the activity of ATP hydrolase. She used shortening of strips of muscle tissue caused by contraction as evidence that ATP was being hydrolysed.

- She took four slides **A**, **B**, **C** and **D**, and added strips of muscle tissue of the same length to each slide.
- She then added the same volume of ATP solutions of different concentrations to the four slides and left each slide for five minutes.
- She then recorded the final length of each strip of muscle tissue.

Her results can be seen in Table 1.

Slide	Concentration of ATP solution added to slide / ×10 ⁻⁶ mol dm ⁻³	Final length of muscle tissue after 5 minutes / mm
A	2	36
В	4	31
С	6	29
D	8	26

Table 1

Other than those given, name **two** variables the student should have controlled.

[2 marks]

0 8 . 3





0 8 4	Describe and explain the pattern shown by the data in Table 1 . [2 marks]	Do not write outside the box
	Description	
	Explanation	
0 8 5	The hydrolysis of 1 dm ³ of a 1 mol dm ^{-3} solution of ATP releases 30 500 J of energy.	
	60% of the energy released during the hydrolysis of 1 mol dm ^{-3} of ATP is released as heat; the rest is used for muscle contraction.	
	The student added 0.05 cm^3 of ATP solution to slide D .	
	Calculate the energy available from ATP for contraction of the muscle on this slide. [3 marks]	
	Answer = J	10
	Turn over ►	




IB/M/Jun18/7401/1

Do not write outside the box

0 9.1 Describe the pathway taken by an oxygen molecule from an alveolus to the blood. [2 marks] 09. 2 Explain how one feature of an alveolus allows efficient gas exchange to occur. [2 marks]








IB/M/Jun18/7401/1

Do not write

box


Carbon monoxide is a poisonous gas that is present in cigarette smoke. This carbon monoxide can be absorbed into the blood where it binds with haemoglobin.

Scientists investigated the concentration of carbon monoxide in cars in which people were smoking or not smoking. They measured the concentration with the car windows open and closed. **Figure 7** shows the scientists' results as they presented them. A value of \pm 2 standard deviations from the mean includes over 95% of the data.





box





Turn over ►

26

Read the following passage.

1 0

Sizes of populations of normal intestinal bacteria are usually controlled by T cells that are produced slowly and in small numbers by the immune system. These T cells do not normally survive for very long. As a result, they do not release large amounts of cytokines. Cytokines are chemicals that can cause swelling of the lining of the intestines.

Crohn's disease is a long-lasting disease that causes swelling of the lining of the intestines. It is believed that Crohn's disease can be caused by a loss of tolerance to normal intestinal bacteria, as shown by an unusually large response by T cells. This response can be triggered by pathogenic bacteria in the intestines of people with a genetic tendency to Crohn's disease.

Some people's Crohn's disease can be controlled by a drug called 5-aminosalicylic acid (5-ASA) that reduces swelling. Another drug called 6-mercaptopurine (6-MP) may also be used. 6-MP inhibits an enzyme required to make adenine and guanine. This is effective because most cells can recycle nucleotides, but T cells are not able to do so.

Use information from the passage and your own knowledge to answer the questions.



10



10.1	The Crohn's disease symptom of swelling of the lining of the intestines could be triggered by pathogenic bacteria in the intestines (lines 6–10).	Do not write outside the box
	Suggest how. [3 marks]	
	Question 10 continues on the next page	



Turn over ►


Do not write outside the box

10.2	Suggest the meaning of 'a genetic tendency to Crohn's disease' (line 10).	[2 marks]
10.3	Suggest why 5 ASA is only offective in controlling the swelling of the lining	of the
	Suggest why 5-ASA is only effective in controlling the swelling of the lining intestines in some people with Crohn's disease (lines 11–12).	[2 marks]




1 0.4	Suggest why 6-MP can be used to control the symptoms of Crohn's disease (lines 13–15).	Do not write outside the box
	Do not include details of enzyme inhibition or protein synthesis in your answer. [3 marks]	
		10
	END OF QUESTIONS	














Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

