ww.xtrapapers.co

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

# **CHEMISTRY**



Paper 5 Practical Test

0620/05

October/November 2005

Candidates answer on the Question Paper.

Additional Materials: As listed in Instructions

to Supervisors

1 hour 15 minutes

Candidate Name							
Centre Number				Candidate Number			

#### **READ THESE INSTRUCTIONS FIRST**

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen in the spaces provided on the Question Paper.

You may use a pencil for any diagrams, graphs or rough working.

DO NOT WRITE IN THE BARCODE.

DO NOT WRITE IN THE GREY AREAS BETWEEN THE PAGES.

Do not use staples, paper clips, highlighters, glue or correction fluid.

You may use a calculator.

Answer all questions.

The number of marks is given in brackets [ ] at the end of each question or part question.

Practical notes are provided on page 8.

FOR EXAMINER'S USE		
1		
2		
Total		

This document consists of 7 printed pages and 1 blank page.

1 You are going to investigate the speed of reaction between aqueous hydrochloric a marble chips (calcium carbonate).

Read **all** the **Instructions** below carefully before starting the Experiments.

#### Instructions

Put 5 test-tubes in a line in the rack provided so you can see the graph paper through them.

To each test-tube you are going to add 3 cm<sup>3</sup> of different solutions of aqueous hydrochloric acid and a marble chip. The marble chips are the same size.

#### Experiment 1

Using the measuring cylinder pour 3 cm<sup>3</sup> of the solution **P** of aqueous hydrochloric acid into the first test-tube.

# **Experiment 2**

Using the measuring cylinder pour 3 cm<sup>3</sup> of the solution **Q** of aqueous hydrochloric acid into the second test-tube.

#### Experiments 3, 4 and 5

Repeat Experiment 1 using 3 cm<sup>3</sup> of the solutions of aqueous hydrochloric acid **R**, **S** and **T** in the third, fourth and fifth test-tubes.

Into all of the test-tubes quickly place a marble chip and start the timer. Shake the tubes from time to time.

Look at the tubes from the side. Take the time in seconds for **each** tube when the lines on the graph paper can be seen through all of the acid in that tube. **Do not stop the timer until all the reactions are finished.** 

Record the times in the table.


# Table of results

Experiment	solution of hydrochloric acid	time/s
1	Р	
2	Q	
3	R	
4	s	
5	Т	

(a) Plot your results on the grid. Draw a best-fit straight line graph.



time/s



[4]

(b)		scribe how the appearance of the mixture in the test-tubes changed as you timed to	the
			[3]
(c)	(i)	Which Experiment has the fastest rate of reaction?	[1]
	(ii)	Explain why this Experiment has the fastest rate.	
			••••
			101

(d)	(i)	In the Experiments which of the reactants is in excess?	For iner's
	(ii)	Explain your answer to (d)(i).	For iner's
			[2]
(e)	(i)	State two sources of error in the Experiments.	
		2	[2]
	(ii)	Suggest two improvements to reduce the sources of error in the Experiments.  1	
		2	
			[2]

		* www.xt	rapapers.com
		5	
2	You are provided with a solid compound	X.	For siner's
	Carry out the following tests on <b>X</b> , recowrite any conclusions in the table.	rding all of your observations in the table. Do	For viner's
	tests	observations	OH
	(a) Appearance of solid X.		
			[2]
	(b) Place one spatula measure of X into a hard-glass test-tube. Heat gently then strongly. Test any gases with damp pH indicator paper. Note all observations.		
			[4]
	Add the rest of solid <b>X</b> to a test-tube. Add about 10 cm <sup>3</sup> of distilled water and shake to dissolve. Divide the solution into five portions in test-tubes.		
	(c) (i) By using a teat pipette add drops of aqueous sodium hydroxide to the first portion of the solution. Now add excess aqueous sodium hydroxide to the		
	test-tube.		[3]
			[2]
	(ii) Using the second portion repeat Experiment (c)(i) using aqueous ammonia instead of aqueous sodium hydroxide.		
			[3]
		•	—— I

		6 Www.xtra	papers.com
	(iii) To the third portion of solution add a few drops of hydrochloric acid and about 1 cm <sup>3</sup> of barium chloride solution.	[1]	Papers.com For iner's
	<ul> <li>(iv) To the fourth portion of solution add a few drops of nitric acid and about 1 cm<sup>3</sup> of lead nitrate solution.</li> </ul>		
		[1]	1
	(v) To the fifth portion of solution add aqueous sodium hydroxide and a spatula measure of aluminium granules.		
	Warm <b>carefully</b> and test the gas with damp indicator paper.	[2	1
(d)	What do tooto (a)(iii) and (iv) tall you	u about <b>V</b> 2	
(u)	What do tests (c)(iii) and (iv) tell you	a about <b>x</b> ?	11
(e)	What conclusions can you draw abo		
		[c	21

7

**BLANK PAGE** 

## **NOTES FOR USE IN QUALITATIVE ANALYSIS**

## **Test for anions**

Test for anions	8 NOTES FOR USE IN QUALITATIVE	ANALYSIS  test result  effervescence, carbon dioxide
anion	test	test result
carbonate (CO 3 <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I⁻) [in solution]	acidify with dilute nitric acid, then aqueous lead(II) nitrate	yellow ppt.
nitrate (NO <sub>3</sub> ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulphate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

# Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia	
aluminium (Al 3+)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess	
ammonium (NH <sub>4</sub> <sup>+</sup> )	ammonia produced on warming	-	
calcium (Ca <sup>2+</sup> )	white., insoluble in excess	no ppt., or very slight white ppt.	
copper(Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution	
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess	
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess	
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution	

# **Test for gases**

gas	test and test results
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	"pops" with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.