

Write your, Centre number, candidate number and name on all the work you hand in.Write in dark blue or black pen.You may use a pencil for any diagrams, graphs or rough working.Do not use staples, paper clips, highlighters, glue or correction fluid.DO NOT WRITE IN ANY BARCODES

Answer **all** questions. Practical notes are provided on page 8.

At the end of the examination, fasten all you work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
Total		

This document consists of **7** printed pages and **1** blank page.



www.papaCambridge.com 1 You are going to investigate the addition of four different solids, A, B, C and D, to 4 g of each solid will be used.

Read **all** the instructions below carefully **before** starting the experiments.

Instructions

Experiment 1

By using a measuring cylinder, pour 30 cm³ of distilled water into one of the polystyrene cups provided. Measure the initial temperature of the water and record it in the table below. Add solid **A** to the water in the cup and stir the mixture with the thermometer. Record the temperature reached after 2 minutes.

Remove the thermometer and rinse with water.

Experiment 2

Repeat Experiment 1 using solid **B** instead of solid **A** and a clean polystyrene cup. Record the initial and final temperatures in the table. Keep the solution for Experiment 5.

Experiment 3

Repeat Experiment 1, using solid **C** and a clean polystyrene cup. Record the temperatures in the table.

Experiment 4

Repeat Experiment 1 using solid **D** and a clean polystyrene cup. Record the temperatures in the table.

Experiment 5


Pour about 2 cm³ of the solution from Experiment 2 into a test-tube. By using a teat pipette add a little of the solution from Experiment 4 to the test-tube. Record your observations.

[2]

Table of results

experiment	initial temperature/°C	final temperature/°C	difference/°C
1			
2			
3			
4			

[5]



	rapapers.con
4	Camp For iner's
(d) Predict the temperature change that would happen if	Can For
(i) 8g of solid B were used in Experiment 2,	[1] The co
	[1] ""."Con
(ii) 60 cm ³ of water was used in Experiment 4.	12
	[1]
(iii) Explain your answer to (d)(ii).	
	[2]
(e) Suggest an explanation for the observations to Experiment 5.	
	[2]
[Total:	20]

2 You are provided with two solutions K and L, each containing the same anion but metal cations.

ation ut th	viae		230
	s.		, each containing the same anion but a
		lowing tests on the solutions, conclusions in the table.	recording all of your observations in the table
		tests	5 , each containing the same anion but of recording all of your observations in the table observations
(a)	Des	cribe the appearance of the	Observations
()		itions.	
	solu	ition K	
	solu	ition L	[1]
(b)	Usir	ng Universal Indicator paper	
	test	the pH of each solution.	
	solu	ition K	рН
	solu	ition L	pH[1]
s on	solu	ition K	
(c)	(i)	By using a teat pipette add drops of aqueous sodium hydroxide to about 3 cm ³ of solution K . Now add excess aqueous sodium	
		hydroxide to the test-tube.	[2]
	(ii)	Repeat experiment (c)(i) using aqueous ammonia instead of aqueous sodium hydroxide.	
			[2]
	(iii)	To about 3 cm ³ of solution K add a few drops of hydrochloric acid and about	
		hydrochloric acid and about 1 cm ³ of barium chloride solution.	[1]

tests	observations
 (iv) To about 3 cm³ of soluti K add a few drops of ni acid and about 1 cm³ of silver nitrate solution. 	tric
ests on solution L	
(d) (i) Repeat experiment (c)(using about 3 cm ³ of solution L.	i) [2]
(ii) Repeat experiment (c)(using about 3 cm ³ of solution L.	ii) [1]
(iii) Repeat experiment (c)(using about 3 cm ³ of solution L.	iii) [1]
(iv) Repeat experiment (c)(using about 3 cm ³ of solution L.	iv) [2]
What does test (b) indicate?	[1]
dentify the anion present in solutior	is K and L .
dentify the metal cation present in	
(i) solution K ,	
	[1]
ii) solution L.	[2]



BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of

NOTES FOR USE IN QUALITATIVE ANALYSIS

Test for anions

8 NOTES FOR USE IN QUALITATIVE ANALYSIS Test for anions anion test			
anion	test	test result	
carbonate (CO ₃ ^{2–})	add dilute acid	effervescence, carbon dioxide produced	
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.	
iodide (I [−]) [in solution]	acidify with dilute nitric acid, then aqueous lead(II) nitrate	yellow ppt.	
nitrate (NO $\overline{3}$) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced	
sulphate (SO ₄ ^{2–}) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.	

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (A <i>l</i> ³⁺)	white ppt., soluble in excess giving a colourless solution white ppt., insoluble in excess	
ammonium (NH ₄ ⁺)	ammonia produced on warming	-
calcium (Ca ²⁺)	white., insoluble in excess	no ppt., or very slight white ppt.
copper(Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Test for gases

gas	test and test results
ammonia (NH ₃)	turns damp red litmus paper blue
carbon dioxide (CO ₂)	turns limewater milky
chlorine (Cl ₂)	bleaches damp litmus paper
hydrogen (H ₂)	"pops" with a lighted splint
oxygen (O ₂)	relights a glowing splint