

Candidates answer on the Question Paper.

No additional materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.


| For Exam | iner's Use |
|----------|------------|
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| Total | |

This document consists of 12 printed pages.





very cle A steel spoon can be coated in silver using electrolysis. The spoon must be very clear 2 free of grease.



(a) Suggest

(i) one advantage of putting a thin layer of silver on the spoon,

| | | . [1] |
|-----|--|--------|
| | (ii) one disadvantage if the spoon is used frequently, | |
| | | [1] |
| | (iii) why the spoon must be very clean and free of grease? | |
| | | . [1] |
| (b) | Which electrode should be the spoon? | |
| | | . [1] |
| (c) | Identify the metal from which the other electrode is made. | |
| | | . [1] |
| | [Tot | al: 5] |

| | | May Waxt | Competence |
|---|-----|---|------------|
| | | 4 | |
| 3 | Thr | ee unlabelled bottles of chemicals each contained one of the following liquids: | For For |
| | • | sodium nitrate dissolved in water; | "Dri |
| | • | pure water; | 30 |
| | • | hexene. | |
| | (a) | Give a test by which you could identify sodium nitrate solution. | |
| | | test | |
| | | result | [2] |
| | | | |
| | (b) | Give a test by which you could identify pure water. | |
| | | test | |
| | | result | [2] |
| | | | |
| | (c) | Give a test by which you could identify hexene. | |
| | | test | |
| | | result | [2] |
| | | | |
| | | [Total: | 6] |

www.papaCambridge.com A student investigated the temperature change produced when equal leng magnesium ribbon reacted with excess dilute sulfuric acid of different concentration. 4 (labelled solutions A, B, C, D and E).

Five experiments were carried out.

Experiment 1

Using a measuring cylinder, 20 cm³ of sulfuric acid solution **A** was poured into a beaker. The initial temperature of the solution was measured. A length of magnesium ribbon was added to the solution and stirred. The highest temperature reached was measured.

Experiment 2

Experiment 1 was repeated using solution **B** instead of solution **A**. The initial and highest temperatures were measured as before.

Experiment 3

Experiment 1 was repeated using solution C. The initial and highest temperatures were measured.

Experiment 4

Experiment 1 was repeated using solution **D**. The initial and highest temperatures were measured.

Experiment 5

Experiment 1 was repeated using solution E. The initial and highest temperatures were measured.

Use the thermometer diagrams in the table on page 6, to record the initial and highest temperatures in each experiment.

| | | | 6 | | town, par | For iner's |
|---------------------------|------------------------|-------------------------------|------------------------|-------------------------------|---------------------------------|---------------|
| solution of sulfuric acid | thermometer diagram | initial temperature /°C | thermometer diagram | highest temperature /°C | change in temperature /°C | iner's |
| A | 30 -25 -20 | | | | | Som |
| в | 30 -25 -20 | | -40 -35 -30 | | | |
| с | 25 20 | | 40 -35 -30 | | | |
| D | 30 -25 -20 | | 35 30 -25 | | | |
| E | 25 20 | | -30 -25 -20 | | | |

[4]

(a) Work out the temperature change for each experiment and record the values in the table. [1]

- 7 (b) Draw a labelled bar chart of the results for Experiments 1, 2, 3, 4 and 5 on the gn below. change in temperature /°C Use the results and observations to answer the following questions. (c) What type of chemical reaction occurs when magnesium reacts with dilute sulfuric
- (d) (i) Which Experiment produced the largest temperature change?


 [1]
 (ii) Suggest why this Experiment produced the largest temperature change.

 [1]
 [1]


acid?

| WWW Xti | |
|--|------|
| 8 (e) Predict the effect on the temperature changes that would happen if (i) equal masses of magnesium powder were used in the Experiments | |
| (e) Predict the effect on the temperature changes that would happen if | Can |
| (i) equal masses of magnesium powder were used in the Experiments. | .101 |
| | [1] |
| (ii) 40 cm ³ of dilute sulfuric acid was used in Experiment 1. | |
| | [1] |
| (iii) Explain your answer to (e)(ii). | |
| | |
| | [1] |
| (f) Give one possible source of experimental error in this investigation. | |
| | |
| | [1] |
| [Total:] | 161 |
| [rotal. | |

| | | www.xtrap |
|-----|---|---|
| Th | e tests on the solutions and some o | 9 were analysed. L was a solution of sodium hydrof the observations are in the table. e. Do not write any conclusions in the table. observations |
| | tests | observations |
| (a) | Appearance of the solutions. | |
| | solution K | colourless liquid |
| | solution L | colourless liquid |
| | solution M | colourless liquid |
| (b) | Universal Indicator paper was used to test the pH of each solution. | |
| | solution K | рН 10 |
| | solution L | pH [1] |
| | solution M | pH 2 |
| (c) | tests on solution K | |
| (| i) Drops of solution K were added to copper sulfate solution in a test-tube. Excess of solution K was then added to the test-tube. | pale blue precipitate formed deep blue solution formed |
| (i | i) Experiment (c)(i) was repeated using aqueous aluminium sulfate instead of aqueous copper sulfate. | white precipitate formed insoluble in excess |
| (ii | i) A few drops of nitric acid and silver nitrate solution were added to solution K. | no visible reaction |
| (d) | tests on solution L | |
| (| i) Experiment (c)(i) was repeated using solution L. | [1] |
| (i | i) Experiment (c)(ii) was repeated using solution L. | [3] |
| (e) | test on solution M | |
| | Experiment (c)(iii) was repeated using solution M . | white precipitate formed |




6 Dilute hydrochloric acid was added to excess calcium carbonate in a beaker as shown.



The beaker was placed on a balance and the mass of the beaker and contents recorded every minute.

The results are shown in the table.

| mass of beaker and contents/g | 184.0 | 178.0 | 175.6 | 174.6 | 174.0 | 174.0 |
|----------------------------------|-------|-------|-------|-------|-------|-------|
| time/min | 0 | 1 | 2 | 3 | 4 | 5 |



| | www.xtra | apa |
|-----|--|-----|
| | 12 | |
| | 12 ves from trees contain a mixture of coloured pigments which are not soluble in water dent was given these two instructions to investigate the pigments in the leaves. Crush some leaves to extract the coloured pigments. Use the liquid extract to find the number of coloured pigments in the leaves. | an |
| | Crush some leaves to extract the coloured pigments. Jse the liquid extract to find the number of coloured pigments in the leaves. | |
| (a) | What would the student need in order to effectively carry out instruction 1? | |
| | | |
| | | |
| | | |
| | [| 3] |
| | | |
| (b) | Describe an experiment to carry out instruction 2. A space has been left below if you want to draw a diagram to help answer the question | n. |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | ••• |
| | | ••• |
| | | |
| | | |
| | | 4] |
| | [Total:] | /] |
| | | |

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.