



**Cambridge International Examinations**  
Cambridge International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 6 1 9 4 7 1 1 3 0 1 \*



**CHEMISTRY**

**0620/52**

Paper 5 Practical Test

**May/June 2015**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

**For Examiner's Use**

**Total**

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.


This document consists of **8** printed pages and **1** insert.

- 1 You are going to investigate the rate of reaction between hydrochloric acid and aqueous sodium thiosulfate. When these chemicals react they form a precipitate which makes the solution go cloudy. The formation of this precipitate can be used to show how fast the reaction proceeds.

**Read all the instructions below carefully before starting the experiments.**

### Instructions

You are going to carry out five experiments using the apparatus shown below.



#### (a) Experiment 1

Use the large measuring cylinder to pour 50 cm<sup>3</sup> of aqueous sodium thiosulfate into a conical flask. Place the conical flask on the printed insert provided.

Fill the small measuring cylinder with 10 cm<sup>3</sup> of the hydrochloric acid provided.

Add the acid to the solution in the conical flask and immediately start your timer and swirl the mixture.

Measure the time taken for the printed text to disappear from view. Record the time in the table. Pour the solution away and rinse the conical flask with distilled water.

#### (b) Experiment 2

Use the large measuring cylinder to pour 40 cm<sup>3</sup> of aqueous sodium thiosulfate into the conical flask, followed by 10 cm<sup>3</sup> of distilled water. Place the conical flask on the printed insert.

Fill the small measuring cylinder with 10 cm<sup>3</sup> of the hydrochloric acid provided.

Add the acid to the solution in the flask, start your timer and swirl the mixture.

Measure the time taken for the printed text to disappear from view. Record the time in the table.

**(c) Experiment 3**

Repeat Experiment 2 using 35 cm<sup>3</sup> of aqueous sodium thiosulfate and 15 cm<sup>3</sup> of distilled water. Record the time in the table.

**(d) Experiment 4**

Repeat Experiment 2 using 30 cm<sup>3</sup> of aqueous sodium thiosulfate and 20 cm<sup>3</sup> of distilled water. Record the time in the table.

**(e) Experiment 5**

Repeat Experiment 2 using 20 cm<sup>3</sup> of aqueous sodium thiosulfate and 30 cm<sup>3</sup> of distilled water. Record the time in the table.


**(f) Complete the table.**

Experiment	volume of aqueous sodium thiosulfate/cm <sup>3</sup>	volume of distilled water/cm <sup>3</sup>	time for printed text to disappear/s
1			
2			
3			
4			
5			

[4]

(g) Plot the results you have obtained on the grid and draw a smooth line graph.

time for printed  
text to disappear / s



[4]

(h) Describe the appearance of the solution in the conical flask at the end of each experiment.

..... [1]

(i) (i) **From your graph**, deduce the time for the printed text to disappear if the experiment was repeated using 25 cm<sup>3</sup> of aqueous sodium thiosulfate and 25 cm<sup>3</sup> of distilled water. Show clearly **on the grid** how you worked out your answer.

..... [3]

(ii) Sketch **on the grid** the curve you would expect if the experiments were repeated at a lower temperature. Label this curve. [1]

(j) (i) In which experiment was the rate of reaction greatest?

..... [1]

(ii) Explain why the rate of reaction was greatest in this experiment.

.....

.....

..... [1]

(k) A student carried out a sixth experiment using 60 cm<sup>3</sup> of aqueous sodium thiosulfate.

Why would this not be an appropriate volume to use in this series of experiments?

.....

..... [2]

(l) Suggest and explain the effect of

(i) using a burette to measure the volume of the hydrochloric acid,

.....

..... [2]

(ii) using a 100 cm<sup>3</sup> conical flask.

.....

..... [2]

[Total: 21]

- 2 You are provided with a mixture of two solids, **J** and **K**, which are both salts. **J** is water soluble and **K** is insoluble.

Carry out the following tests on the mixture, recording all of your observations in the table. Conclusions must **not** be written in the table.

tests	observations
<b>(a)</b> Describe the appearance of the mixture.	..... [1]
<p>Add 10 cm<sup>3</sup> of distilled water to the mixture. Stopper the boiling tube and shake the mixture for a minute. Filter the mixture and keep the residue for tests later.</p> <p><u>tests on the filtrate</u></p> <p>Divide the solution into three equal portions in three test-tubes.</p> <p><b>(b)</b> To the first portion of the solution, add about 1 cm<sup>3</sup> of aqueous sodium hydroxide. Heat the mixture gently and test any gas evolved with damp pH indicator paper.</p>	<p>.....</p> <p>..... [3]</p>
<b>(c)</b> To the second portion of the solution, add a few drops of dilute nitric acid and about 1 cm <sup>3</sup> of aqueous silver nitrate solution.	..... [2]
<p><b>(d)</b> To the third portion of the solution, add about 1 cm<sup>3</sup> of hydrogen peroxide solution.</p> <p>Add a spatula measure of starch to the mixture.</p> <p>Test the gas given off with a splint.</p>	<p>.....</p> <p>.....</p> <p>..... [5]</p>
<p><u>tests on the residue</u></p> <p>Use a spatula to transfer some of the residue into a test-tube.</p> <p><b>(e)</b> Add about 3 cm<sup>3</sup> of dilute hydrochloric acid to the test-tube. Test the gas given off.</p> <p>To the solution add about 1 cm<sup>3</sup> of dilute sulfuric acid.</p>	<p>.....</p> <p>..... [3]</p> <p>..... [1]</p>

(f) What conclusions can you draw about solid J?

.....  
..... [2]

(g) What conclusions can you draw about solid K?

.....  
.....  
..... [2]

[Total: 19]

## NOTES FOR USE IN QUALITATIVE ANALYSIS

## Test for anions

<i>anion</i>	<i>test</i>	<i>test result</i>
carbonate ( $\text{CO}_3^{2-}$ )	add dilute acid	effervescence, carbon dioxide produced
chloride ( $\text{Cl}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide ( $\text{I}^-$ ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate ( $\text{NO}_3^-$ ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate ( $\text{SO}_4^{2-}$ ) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

## Test for aqueous cations

<i>cation</i>	<i>effect of aqueous sodium hydroxide</i>	<i>effect of aqueous ammonia</i>
aluminium ( $\text{Al}^{3+}$ )	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium ( $\text{NH}_4^+$ )	ammonia produced on warming	–
calcium ( $\text{Ca}^{2+}$ )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper ( $\text{Cu}^{2+}$ )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) ( $\text{Fe}^{2+}$ )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) ( $\text{Fe}^{3+}$ )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc ( $\text{Zn}^{2+}$ )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

## Test for gases

<i>gas</i>	<i>test and test results</i>
ammonia ( $\text{NH}_3$ )	turns damp red litmus paper blue
carbon dioxide ( $\text{CO}_2$ )	turns limewater milky
chlorine ( $\text{Cl}_2$ )	bleaches damp litmus paper
hydrogen ( $\text{H}_2$ )	'pops' with a lighted splint
oxygen ( $\text{O}_2$ )	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.