



## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER	CANDIDATE NUMBER	

CHEMISTRY 0620/51

Paper 5 Practical Test October/November 2016

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Practical notes are provided on pages 11 and 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use
Total

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 10 printed pages and 2 blank pages.



1 You are going to investigate what happens when nitric acid reacts with aqueous solutions of two different alkalis, solution **N** and solution **O**.

#### Read all the instructions carefully before starting the experiments.

#### Instructions

You are going to carry out two experiments.

## (a) Experiment 1

Use a measuring cylinder to pour 50 cm<sup>3</sup> of solution **N** into the polystyrene cup provided. Put the polystyrene cup into a 250 cm<sup>3</sup> beaker for support. Measure the initial temperature of the solution and record it in the first row of the table.

Fill the burette with nitric acid to the 0.0 cm<sup>3</sup> mark.

Add 5.0 cm<sup>3</sup> of nitric acid to solution **N** in the polystyrene cup and stir the solution.

Measure and record the maximum temperature of the solution in the table.

Add a further 5.0 cm<sup>3</sup> of nitric acid to the polystyrene cup and stir the solution. Measure and record the maximum temperature of the solution in the table.

Continue to add 5.0 cm<sup>3</sup> portions of nitric acid to the polystyrene cup, until a total volume of 40 cm<sup>3</sup> of nitric acid has been added. Stir after each addition and measure and record the maximum temperatures in the table.

Pour the solution away and rinse the polystyrene cup.

volume of nitric acid added/cm <sup>3</sup>	maximum temperature of the solution in the polystyrene cup/°C
0.0	
5.0	
10.0	
15.0	
20.0	
25.0	
30.0	
35.0	
40.0	


# (b) Experiment 2

Refill the burette with nitric acid. Repeat Experiment 1 using solution  $\bf 0$  instead of solution  $\bf N$ . Record your results in the table.

volume of nitric acid added/cm³	maximum temperature of the solution in the polystyrene cup/°C
0.0	
5.0	
10.0	
15.0	
20.0	
25.0	
30.0	
35.0	
40.0	

[2]

(c) Plot the results for Experiments 1 and 2 on the grid and draw **two** smooth line graphs. Clearly label your graphs.



(d) Use your graph to estimate the maximum temperature of the solution when 13 cm³ of nitric acid were added to 50 cm³ of solution N in Experiment 1.

Show clearly **on the grid** how you worked out your answer.

	00	[2]
 	$\cup$	[4]

(e)	Name a suitable indicator that could be used in Experiment 1.
	[1]
(f)	Solution <b>N</b> and solution <b>O</b> are the same concentration.
	In which experiment is the temperature change greater? Suggest why the temperature change is greater in this experiment.
	[2]
(g)	How would the results differ in Experiment 1 if 100 cm³ of solution <b>N</b> were used?
	[1]
(h)	Suggest why a polystyrene cup is used in these experiments and <b>not</b> a copper can.
	[1]
(i)	State <b>one</b> source of error in the experiments. Suggest an improvement to reduce this source of error.
	source of error
	improvement[2]
	[2]

[Total: 17]

2	You are provided with solid <b>P</b> , which is a metallic salt.
	Carry out the following tests on solid P, recording all of your observations at each stage.

tests on solid P			
(a) De	escribe the appearance of solid <b>P</b> .		
	[1]		
<b>(b)</b> Us	e a spatula to divide solid <b>P</b> into three portions.		
(i)	test 1 Heat the first portion of solid <b>P</b> in a hard-glass test-tube. Test any gases given off with cobalt(II) chloride paper. Record your observations.		
	[3]		
(ii)	test 2 Carry out a flame test on the second portion of solid <b>P</b> . Record your observations.		
	[1]		
tests o	on a solution of P		
	out $10\text{cm}^3$ of distilled water to the third portion of solid <b>P</b> in a test-tube. Stopper the test-tube ake it to dissolve solid <b>P</b> .		
(c) Div	vide the solution into four equal portions in four test-tubes. Carry out the following tests.		
(i)	To the first portion of the solution, add several drops of aqueous sodium hydroxide. Then add excess aqueous sodium hydroxide to the mixture. Record your observations.		
	[3]		
(ii)	To the second portion of the solution, add excess aqueous ammonia. Record your observations.		

	(iii)	To the third portion of the solution, add a few drops of dilute nitric acid and about 1 cm <sup>3</sup> of aqueous silver nitrate.  Record your observations.	of
		[	1]
	(iv)	To the fourth portion of the solution, add a few drops of dilute nitric acid and about 1 cm <sup>3</sup> of aqueous barium nitrate.  Record your observations.	
		[2	
(d)	Sol	lid <b>P</b> contains a metal ion.	
	Su	ggest what the appearance of solid <b>P</b> in <b>(a)</b> tells you about the identity of the metal ion.	
		[	1]
(e)	Wh	nat does <b>test 1</b> tell you about solid <b>P</b> ?	4.
		[	1]
(f)	Wh	nat does test 2 tell you about solid P?	
		[	1]
(g)	lde	ntify solid <b>P</b> .	
		[/	
		[Total: 17	_
		L. C.	- 4

3

lds
id.
[6]
6]

9

# **BLANK PAGE**

10

# **BLANK PAGE**

# NOTES FOR USE IN QUALITATIVE ANALYSIS Test for anions

anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide (Br <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO <sub>3</sub> <sup>-</sup> ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify, then add aqueous barium nitrate	white ppt.
sulfite (SO <sub>3</sub> <sup>2-</sup> )	add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide	sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless

# **Test for aqueous cations**

	T	1
cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+) white ppt., soluble in excess giving a colourless solution		white ppt., insoluble in excess
ammonium (NH <sub>4</sub> <sup>+</sup> ) ammonia produced on warming		-
calcium (Ca <sup>2+</sup> )	white ppt., insoluble in excess	no ppt. or very slight white ppt.
chromium(III) (Cr³+) green ppt., soluble in excess		grey-green ppt., insoluble in excess
copper (Cu²+) light blue ppt., insoluble in excess		light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe³+) red-brown ppt., insoluble in excess		red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess, giving a colourless solution

### **Test for gases**

gas	test and test results	
ammonia (NH <sub>3</sub> )	turns damp, red litmus paper blue	
carbon dioxide (CO <sub>2</sub> )	turns limewater milky	
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper	
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint	
oxygen (O <sub>2</sub> )	relights a glowing splint	
sulfur dioxide (SO <sub>2</sub> )	turns acidified aqueous potassium manganate(VII) from purple to colourless	

### Flame tests for metal ions

metal ion	flame colour
lithium (Li <sup>+</sup> )	red
sodium (Na <sup>+</sup> )	yellow
potassium (K+)	lilac
copper(II) (Cu <sup>2+</sup> )	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.