

CANDIDATE

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

White Con

*	
9	
σ	
N	
W	
N	
œ	
9	
W	
_	
7	

NAME			 	
CENTRE NUMBER		CANDIDATE NUMBER		
-				

MATHEMATICS

0580/21

Paper 2 (Extended)

May/June 2011

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials:

Electronic calculator

Mathematical tables (optional)

Geometrical instruments

Tracing paper (optional)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

If working is needed for any question it must be shown below that question.

Electronic calculators should be used.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place.


For π , use either your calculator value or 3.142.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.


The total of the marks for this paper is 70.

Answer [1]





 $A \cup B'$



 $(A \cap B)'$

[2]

 $81^{0.25} \div 4^{-2}$. Calculate 3

> Answer [2]

(a) Find m when $4^m \times 4^2 = 4^{12}$.

$$Answer(a) m =$$
 [1]

(b) Find *p* when $6^p \div 6^5 = \sqrt{6}$.

$$Answer(b) p =$$
 [1]

WW. Pak		pers.com
	aCanne	For iner's
	[1]	COM

- 5 A hummingbird beats its wings 24 times per second.
 - (a) Calculate the number of times the hummingbird beats its wings in one hour.


1	Г17
Answer(a)	111
11.00 (00)	 L + J

(b) Write your answer to **part (a)** in standard form.

Answer(b)	[1]
-----------	-----

6





NOT TO SCALE

A company makes solid chocolate eggs and their shapes are mathematically similar.

The diagram shows eggs of height 2 cm and 6 cm.

The mass of the small egg is 4 g.

Calculate the mass of the large egg.


7 Find the length of the straight line from Q(-8, 1) to R(4, 6).

$$Answer\ QR =$$
 [3]

Calculate the radius of a sphere with volume 1260 cm³. 8 [The volume, V, of a sphere with radius r is $V = \frac{4}{3}\pi r^3$.]

4		F 2 7
Answer	 cm	[3]

9



NOT TO **SCALE**

AB is parallel to CD. Calculate the value of *x*.

$$Answer x = [3]$$

10 Solve the simultaneous equations.

$$3x + y = 30$$
$$2x - 3y = 53$$

$$Answer x =$$

$$y =$$
 [3]

			₩ ₩	w.xtrapapers.co
	5			D
A re	ectangular photograph measures 23.3 cm by 19.7 cm culate the lower bound for	, each correct	to 1 decimal place.	For iner's
(a)	the perimeter,			Tage
				1 ag
		Answer(a)		cm [2]
(b)	the area.			
		Answer(b)		cm ² [1]
A tr	rain leaves Barcelona at 21 28 and takes 10 hours an	d 33 minutes	to reach Paris.	
(a)	Calculate the time the next day when the train arriv	ves in Paris.		
		Answer(a)		[1]
(b)	The distance from Barcelona to Paris is 827 km.			
	Calculate the average speed of the train in kilometr	es per hour.		
		Answer(b)		km/h [3]
	(a) A to (a)	A rectangular photograph measures 23.3 cm by 19.7 cm Calculate the lower bound for (a) the perimeter, (b) the area. A train leaves Barcelona at 21 28 and takes 10 hours an (a) Calculate the time the next day when the train arrivation of the content of the co	A rectangular photograph measures 23.3 cm by 19.7 cm, each correct Calculate the lower bound for (a) the perimeter, Answer(a) (b) the area. Answer(b) A train leaves Barcelona at 21 28 and takes 10 hours and 33 minutes (a) Calculate the time the next day when the train arrives in Paris. Answer(a) (b) The distance from Barcelona to Paris is 827 km. Calculate the average speed of the train in kilometres per hour.	A rectangular photograph measures 23.3 cm by 19.7 cm, each correct to 1 decimal place. Calculate the lower bound for (a) the perimeter, Answer(a) Answer(b) A train leaves Barcelona at 21 28 and takes 10 hours and 33 minutes to reach Paris. (a) Calculate the time the next day when the train arrives in Paris. Answer(a) Answer(b) Answer(a) Calculate the average speed of the train in kilometres per hour.

12	The seels on a man	:.	1.	20	000
13	The scale on a map	18	1:	20	UUU.

(a) Calculate the actual distance between two points which are 2.7 cm apart on the map. Give your answer in kilometres.

Answer(a)	km	[2]
Answer (u)	 VIII	141

(b) A field has an area of $64 400 \,\mathrm{m}^2$. Calculate the area of the field on the map in cm².

Answer(b)
$$cm^2$$
 [2]

Solve the equation $2x^2 + 3x - 6 = 0$. Show all your working and give your answers correct to 2 decimal places.


15 A teacher asks 36 students which musical instruments they play.

 $P = \{\text{students who play the piano}\}\$

 $G = \{\text{students who play the guitar}\}\$

 $D = \{ \text{students who play the drums} \}$

The Venn diagram shows the results.



(a) Find the value of x.

$$Answer(a) x =$$
 [1]

(b) A student is chosen at random.

Find the probability that this student

(i) plays the drums but **not** the guitar,

$$Answer(b)(i) \qquad [1]$$


(ii) plays only 2 different instruments.

(c) A student is chosen at random from those who play the guitar.

Find the probability that this student plays no other instrument.

[2]

16



The diagram shows a square of side k cm.

The circle inside the square touches all four sides of the square.


(a) The shaded area is $A ext{ cm}^2$.

Show that $4A = 4k^2 - \pi k^2.$

Answer (a)

(b) Make k the subject of the formula $4A = 4k^2 - \pi k^2$.

Answer(b) k = [3]



A, B and C are points on a circle, centre O.

TA is a tangent to the circle at A and OBT is a straight line.

AC is a diameter and angle $OTA = 24^{\circ}$.

Calculate

(a) angle AOT,


$$Answer(a) \text{ Angle } AOT =$$
 [2]

(b) angle ACB,

$$Answer(b) \text{ Angle } ACB =$$
 [1]

(c) angle ABT.

$$Answer(c) \text{ Angle } ABT =$$
 [2]



In the diagram, PQS, PMR, MXS and QXR are straight lines.

$$PQ = 2 QS$$
.

 \widetilde{M} is the midpoint of PR.

$$QX : XR = 1 : 3.$$

$$\overrightarrow{PQ} = \mathbf{q}$$
 and $\overrightarrow{PR} = \mathbf{r}$.

- (a) Find, in terms of q and r,
 - (i) \overrightarrow{RQ} ,


$$Answer(a)(i) \overrightarrow{RQ} =$$
 [1]

(ii) \overrightarrow{MS} .

$$Answer(a)(ii) \overrightarrow{MS} =$$
 [1]

(b) By finding \overrightarrow{MX} , show that X is the midpoint of MS.

Answer (b)



The diagram shows the speed-time graph of a train journey between two stations. The train accelerates for two minutes, travels at a constant maximum speed, then slows to a stop.

(a) Write down the number of seconds that the train travels at its constant maximum speed.

Answer(a) _____ s [1]

(b) Calculate the distance between the two stations **in metres**.

Answer(b) m [3]

(c) Find the acceleration of the train in the first two minutes.

Give your answer in m/s².

Answer(c) m/s^2 [2]

$$f(x) = x^3$$

$$f(x) = x^3 \qquad g(x) = 2x - 3$$

- (a) Find
 - (i) g(6),

1 ()(i)	Γ1
Answer(a)(1)	- 11
11.00 ,, 0. (00) (1)	 1 -

(ii) f(2x).

(b) Solve fg(x) = 125.

$$Answer(b) x =$$
 [3]

(c) Find the inverse function $g^{-1}(x)$.

$$Answer(c) g^{-1}(x) =$$
 [2]