

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

MARK SCHEME for the May/June 2011 question paper
for the guidance of teachers

0606 ADDITIONAL MATHEMATICS

0606/11

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

Mark Scheme Notes

Marks are of the following three types:

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Accuracy mark for a correct result or statement independent of method marks.
- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
 - The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
 - Note: B2 or A2 means that the candidate can earn 2 or 0.
B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606




The following abbreviations may be used in a mark scheme or used on the scripts:

AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)



Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{\quad}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.
OW –1,2	This is deducted from A or B marks when essential working is omitted.
PA –1	This is deducted from A or B marks in the case of premature approximation.
S –1	Occasionally used for persistent slackness – usually discussed at a meeting.
EX –1	Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>1</p> $\frac{(1 + \cos \theta) + (1 - \cos \theta)}{(1 + \cos \theta)(1 - \cos \theta)}$ $= \frac{2}{1 - \cos^2 \theta}$ $= \frac{2}{\sin^2 \theta}$ $= 2 \operatorname{cosec}^2 \theta$	<p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>M1 for attempt to deal with fractions</p> <p>M1 for attempt at simplification and use of $1 - \cos^2 \theta = \sin^2 \theta$ in denominator</p>
<p>2</p> $\lg ab^3 - \lg 1000$ $= \lg \frac{ab^3}{1000}$	<p>B1B1</p> <p>B1</p> <p>[3]</p>	<p>B1 for $\lg ab^3$, B1 for $\lg 1000$</p>
<p>3 (a) (i)</p>  <p>(ii)</p>  <p>(iii)</p>  <p>(b) $n(P) = 3$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>[4]</p>	<p>B1 for each region shaded correctly</p>
<p>4 (a) Powers of 2: $4(3x - 2) = 3(2x)$ or equivalent for powers of 4, 8 or 16 $x = \frac{4}{3}$, allow 1.33</p> <p>(b) $p = 1, q = -\frac{4}{5}$</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>B1B1</p> <p>[5]</p>	<p>M1 for powers of 2, 4, 8 or 16</p> <p>A1 for all powers correct</p>

Page 5	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>5 (i)</p>  <p>(ii)</p>  <p>(iii) 5</p>	<p>B1 B1 B1</p> <p>√B1</p> <p>√B1 [5]</p>	<p>B1 for shape B1 for 1 cycle between 4 and -2 B1 all correct</p> <p>√B1 for modulus of (i)</p> <p>√ on their graph</p>
<p>6 (i) $3x^2 = -2x^2 + 20x - 20$ and verification.</p> <p>Or $(x - 2)^2 = 0$, $x = 2$</p> <p>(ii) for OA, $\frac{dy}{dx} = 2x$ when $x = 2$, grad = 4 for other curve, $\frac{dy}{dx} = -\frac{4}{3}x + \frac{20}{3}$ when $x = 2$, grad = 4</p> <p>Or $2x = -\frac{4}{3}x + \frac{20}{3}$ leading to $x = 2$</p> <p>(iii) tangent $y - 4 = 4(x - 2)$</p>	<p>B1 B1 B1 M1 A1 M1 M1 A1 B1 [5]</p>	<p>substitution of $x = 2$</p> <p>B1 for solution of equation</p> <p>B1 for grad at A from OA</p> <p>M1 for attempt to differentiate the other curve and substitute $x = 2$</p> <p>M1 for differentiation of both M1 for equating and attempt to solve</p>

Page 6	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>7 Grad of $AB = -2$, perp grad $= \frac{1}{2}$</p> <p>Eqn of perp $y - 15 = \frac{1}{2}(x + 2)$</p> <p>$C(0, 16)$</p> <p>Area $= \frac{1}{2}\sqrt{125}\sqrt{5}$</p> <p>$= 12.5$</p> <p>(or $\frac{1}{2} \begin{vmatrix} -2 & 3 & 0 & -2 \\ 15 & 5 & 16 & 15 \end{vmatrix} = \frac{1}{2}(38 - 13)$)</p>	<p>B1M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>B1 for grad AB</p> <p>M1 use of $m_1 m_2 = -1$</p> <p>M1 for correct attempt to find the equation of AC and hence to find C</p> <p>M1 for a valid method to find area</p>
<p>8 (a) AB, AC</p> <p>(b) Either: $\mathbf{Y} = \mathbf{X} \begin{pmatrix} -12x + 3y & 6 \\ -7x + 3y & 6 \end{pmatrix}$</p> <p>$\begin{pmatrix} 2x & 3y \\ x & 4y \end{pmatrix} = \begin{pmatrix} -52x + 18y & 36 \\ -71x + 24y & 48 \end{pmatrix}$</p> <p>leading to $y = 12$ and $x = 4$</p> <p>Or</p> <p>$\frac{1}{10 - 12} \begin{pmatrix} 5 & -4 \\ -3 & 2 \end{pmatrix} \mathbf{Y} = \begin{pmatrix} -12x + 3y & 6 \\ -7x + 3y & 6 \end{pmatrix}$</p> <p>$-\frac{1}{2} \begin{pmatrix} 6x & -y \\ -4x & -y \end{pmatrix} = \begin{pmatrix} -12x + 3y & 6 \\ -7x + 3y & 6 \end{pmatrix}$</p> <p>leading to $y = 12$ and $x = 4$</p>	<p>B2,1,0</p> <p>M1</p> <p>M1A1</p> <p>M1</p> <p>A1A1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1A1</p> <p>[8]</p>	<p>-1 each one incorrect or extra</p> <p>M1 for pre-multiplying by \mathbf{X}</p> <p>M1 for multiplication of matrices</p> <p>A1 for correct product</p> <p>M1 for equating like elements</p> <p>B1 for determinant for inverse</p> <p>B1 for 'matrix part' of inverse</p> <p>M1 for multiplication of matrices</p> <p>M1 for equating like elements</p>
<p>9 (i) 5</p> <p>(ii) $a = -20 \sin 4t$</p> <p>$\sin 4t = -0.5$</p> <p>$t = \frac{7\pi}{24}$ (allow 0.916)</p> <p>(iii) $s = \frac{5}{4} \sin 4t (+c)$</p> <p>When $t = 5$, $s = 1.14$</p>	<p>B1</p> <p>M1A1</p> <p>DM1</p> <p>A1</p> <p>M1A1</p> <p>DM1</p> <p>A1</p> <p>[9]</p>	<p>M1 for attempt to differentiate</p> <p>DM1 for attempt to solve for $4t$</p> <p>M1 for attempt to integrate</p> <p>DM1 for substitution of t in radians</p>

Page 7	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>10 (a) (i) $2 = a - 3, a = 5$</p> <p>(ii) $y = -5e^{1-x} - x^3 + c$ $c = 10$ $y = -5e^{1-x} - x^3 + 10$</p> <p>(b) (i) $\frac{1}{7} \frac{3}{4} (7x+8)^{\frac{4}{3}}$</p> <p>(ii) $\left[\frac{3}{28} (7x+8)^{\frac{4}{3}} \right]_0^8$ $= \frac{180}{7}$ or 25.7</p>	<p>B1</p> <p>√B1 B1B1 M1 A1</p> <p>B1B1</p> <p>M1A1</p> <p>[10]</p>	<p>√B1 for first term using their a B1 for $-x^3$, B1 for $+c$ M1 for attempt to find c</p> <p>B1 for $\frac{1}{7}$, B1 for $\frac{3}{4} (7x+8)^{\frac{4}{3}}$</p> <p>M1 for use of limits</p>
<p>11 (i) $2(x-2)^2 - 3$</p> <p>(ii) $x \geq 2$ or equivalent</p> <p>(b) (i) $g(x) \geq 4, h^{-1}(x) \geq 0$</p> <p>(ii) Correct sketch</p> <p>(iii) $g(4x-25) = 85$ $(4x-25)^2 + 4 = 85$ $x = \frac{17}{2}, x = 4$ Discarding $x = 4$</p>	<p>B1B1</p> <p>√B1</p> <p>B1B1</p> <p>B1 B1 B1</p> <p>M1 DM1 A1 B1</p> <p>[12]</p>	<p>B1 for -2, B1 for -3</p> <p>√ on their $'-2'$</p> <p>B1 for each</p> <p>B1 for $g(x)$ B1 for $g^{-1}(x)$ B1 for idea of symmetry</p> <p>M1 for correct order DM1 for attempt to solve A1 for both B1 for discarding $x = 4$</p>

Page 8	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>12 EITHER</p> <p>(i) $\frac{dy}{dx} = 3x^2 - 14x + 8$</p> <p>When $\frac{dy}{dx} = 0$, $x = \frac{2}{3}, 4$</p> <p>$\frac{d^2y}{dx^2} = 6x - 14$,</p> <p>$x = \frac{2}{3}$ max, $x = 4$ min</p> <p>(ii) Use of $\frac{dy}{dt} = \frac{dy}{dz} \times \frac{dz}{dt}$, leading to</p> <p>$\frac{dy}{dt} = -\frac{5}{6}$ allow -0.833</p> <p>(iii) Use of $\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$ leading to</p> <p>$\frac{dx}{dt} = \frac{5}{48}$</p>	<p>M1A1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>$\sqrt{A1}$</p> <p>[10]</p>	<p>M1 for attempt to differentiate</p> <p>M1 for attempt to equate to zero and solve</p> <p>M1 for attempt to differentiate (or other valid method)</p> <p>A1 correct from correct working for both</p> <p>M1 for attempt to use rates of change</p> <p>M1 for attempt to use rates of change</p> <p>ft on $\frac{dy}{dt}$</p>
<p>12 OR</p> <p>(i) $2x^2y = 72$, $A = 4x^2 + 6xy$</p> <p>leading to given answer</p> <p>(ii) $\frac{dA}{dx} = 8x - \frac{216}{x^2}$</p> <p>When $\frac{dA}{dx} = 0$, $x = \sqrt[3]{27} = 3$</p> <p>Dimensions are 3 by 6 by 4</p> <p>(iii) Use of $\partial A \approx \frac{dA}{dx} \times \partial x$ leading to</p> <p>$\partial A = -38p$, decrease</p>	<p>B1M1</p> <p>A1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1$\sqrt{A1}$</p> <p>[10]</p>	<p>B1 for $2x^2y = 150$,</p> <p>M1 for $A = 4x^2 + 6xy$</p> <p>M1 for attempt to differentiate</p> <p>M1 for attempt to equate to zero and solve</p> <p>A1 for dimensions</p> <p>M1 for attempt to use small changes</p> <p>A1 for $-38p$, $\sqrt{A1}$ on their ∂A</p>