

Cambridge Assessment International Education Cambridge International General Certificate of Secondary Education (9–1)

#### MATHEMATICS

Paper 6 MARK SCHEME Maximum Mark: 96 0626/06 October/November 2019

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2019 series for most Cambridge IGCSE<sup>™</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

This syllabus is regulated for use in England as a Cambridge International Level 1/Level 2 (9–1) Certificate.

This document consists of 9 printed pages.

[Turn over





### Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit
  is given for valid answers which go beyond the scope of the syllabus and mark scheme,
  referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.



### MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

#### Types of mark

- M Method marks, awarded for a valid method applied to the problem.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation '**dep**' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

#### Abbreviations

answers which round to awrt cao correct answer only dependent dep FT follow through after error isw ignore subsequent working nfww not from wrong working or equivalent oe rounded or truncated rot SC Special Case seen or implied soi



www.xtrapapers.com October/November

2019

Question	Answer	Marks	Partial Marks
1(a)	$\frac{1}{2}(x+x+4)(x-1)$ oe	M1	
	(x+2)(x-1) = 108 or $(2x+4)(x-1) = 216$	M1	
	$x^{2} + 2x - 1x - 2 = 108$ or $2x^{2} + 4x - 2x - 4 = 216$	M1	correct expansion of <i>their</i> brackets, allow one error or omission
	correct completion to $x^2 + x - 110 = 0$	A1	All correct with no errors seen
1(b)	-11 and +10 with correct factorisation seen	3	M2 for $(x+11)(x-10)$ or M1 for $(x+a)(x+b)$ where $ab = -110$ or $a+b=1$ After M1, SC1 for $x = -their a$ and $x = -their b$ If 0 scored, SC1 for both correct answers.
1(c)	9	1	<b>FT</b> provided 2 values of different signs are given for $x$ in <b>(b)</b> .
2(a)	$2x + 5y = 20$ or $y = \frac{-2}{5}x + 4$ or equivalent 3 term equation	3	<b>B1</b> for $-\frac{4}{10}$ <b>M1</b> for $y = \pm \frac{4}{10}x + c$ or, e.g. $(y-4) = \pm \frac{4}{10}(x-0)$
2(b)(i)	21.8 or 21.80	2	<b>M1</b> for $tan[] = \frac{4}{10}$ or $tan[] = \frac{10}{4}$
2(b)(ii)	2.6[0] or 2.599	3	<b>B1</b> for using correct right-angled triangle soi, e.g. marked on diagram. <b>M1FT</b> for $sin(their(b)(i)) = \frac{d}{7}$ oe
3(a)	624	3	M2 for $\frac{642.72}{1.03}$ oe or M1 for 642.72 associated with 103[%]



Question	Answer	Marks	Partial Marks
3(b)	15.8	4	<b>M3</b> for $\frac{1000 \times 1.36}{1000 \times 1.174} \times 100[-100]$ oe soi by 116 or 115.8 or 15.8 or 15.84 or <b>M2</b> for $\frac{1000 \times 1.36}{1.174}$ soi by 1158 or 1160 or <b>M1</b> for 1000 × 1.36
3(c)	6 with supporting evidence	4	M2 for calculating correctly either $400 \times 1.02^n$ or $390 \times 1.025^n$ for $n \ge 3$ oe or M1 if $n = 2$ calculated correctly for either or M1 for $400 \times 1.02^n$ and $390 \times 1.025^n$ both soi M1 for calculating correctly both investments for a value of $n$ , $n \ge 3$ Alternative method: M2 for calculating correctly $\frac{1.025^n}{1.02^n}$ for $n \ge 3$ oe or M1 if $n = 2$ calculated correctly or M1 for $\frac{1.025^n}{1.02^n}$ and $\frac{400}{390}$ both soi M1 for $1.0256[4]$
3(d)(i)	612.49	3	<b>B1</b> for 604 and <b>B1</b> for 608.16
3(d)(ii)	500	1	
	their investment will decrease oe	1	dep on <i>their</i> (d)(ii) $\leq 500$
4(a)	30	2	<b>M1</b> for $\frac{360}{12}$
4(b)(i)	Parallelogram constructed with arcs	2	M1 for one pair of correct intersecting arcs drawn



www.xtrapapers.com October/November 2019

Question	Answer	Marks	Partial Marks
4(b)(ii)	Circle drawn with correct constructions shown	3	<ul> <li>M2 for correct perpendicular bisector for 2 of the other points with correct arcs or M1 for correct perpendicular bisector with incorrect or no arcs or for 2 correct pairs of arcs seen.</li> <li>A1 for correct circle dependent on at least M1</li> <li>If 0 scored SC1 for correct circle</li> </ul>
5(a)(i)		3	<b>B2</b> for three numbers correctly placed or <b>B1</b> for two numbers correctly placed.
5(a)(ii)	12	1	<b>FT</b> 7 + <i>their</i> (4+1)
5(a)(iii)	13	1	
5(b)(i)	$(F \cap G)$ ' oe	1	
5(b)(ii)	$(P \cup Q) \cap R$ oe	1	
6(a)	$y \ge 2 \text{ oe} x + y \ge 9 \text{oe}$	2	B1 for each
6(b)	$400x + 700y \le 5600$ oe	1	
6(c)	4 correct, continuous, ruled lines drawn and correct region left unshaded	6	<ul> <li>B1 for x = 4 drawn</li> <li>B1FT for <i>their</i> y = 2 drawn dep on y = k</li> <li>B1FT for <i>their</i> x + y = 9 drawn dep on <i>their</i> x + y = k oe</li> <li>B2FT for <i>their</i> 4x + 7y = 56 drawn dep on <i>their</i> 4x + 7y = k</li> <li>or SC1FT for line with negative gradient passing through one of <i>their</i> intercepts (14, 0) or (0, 8)</li> </ul>
6(d)	12	1	

Page 6 of 9



	Answer	Marks	Partial Marks
6(e)	5, 5	2	<b>B1</b> for 5,5 or for 27.5[0]
	27.5[0]		If 0 scored <b>SC1</b> for clear use of $[P=]1.5x+4y$ for a point in <i>their</i> region.
7(a)	349 or 349.0 to 349.11	2	<b>M1</b> for $\left[\frac{64}{360}\times\right]\pi\times25^2$
7(b)	48.1 or 48.06	4	<b>B1</b> for 148° or 74° or 16° seen on diagram or used.
			<b>M2</b> for $\frac{25\sin 148}{\sin 16}$ or <b>M1</b> for $\frac{\sin 16}{25} = \frac{\sin 148}{x}$ oe
			OR M2 for $\sqrt{25^2 + 25^2 - 2 \times 25 \times 25 \cos 148}$ or M1 for $25^2 + 25^2 - 2 \times 25 \times 25 \cos 148$
			OR M2 for $2 \times 25 \times \sin 74$ oe or M1 for $\sin 74 = \frac{x}{25}$ oe
7(c)	166 or 165.5 to 165.6	2	M1 for $\frac{1}{2} \times 25 \times 25 \times \sin 148$ or for $\frac{1}{2} \times 25 \times their7(b) \times \sin 16$
7(d)	35.2 or 35.17 to 35.3	4	M3 for their 7(a) + 2×their(c) – $\frac{32}{360} \times \pi \times their 7(b)^{2}$
			or M2 for $\frac{32}{360} \times \pi \times their 7(b)^2$ or for two of the M1 cases
			or M1 for <i>their</i> 7(a) + 2× <i>their</i> (c) or $k \times \pi \times their$ 7(b) <sup>2</sup> or BOC = 32°
8(a)	Two correct limitations with improvement	2	B1 for each
	20, 60, 50, 80	2	B1 for two correct



Question	Answer	Marks	Partial Marks
8(c)	1580	4	<b>M1</b> for midpoints soi (375, 750, 1250, 2000, 3000)
			M1 for use of $\sum fx$ with x in correct interval including both boundaries $20 \times 375 + 60 \times 750 + 50 \times 1250 +$ $80 \times 2000 + 40 \times 3000$
			<b>M1</b> (dep on 2nd M1) for $\sum fx \div 250$
8(d)(i)	Correct graph	4	<b>B1</b> for 6 points plotted at upper ends of intervals
			<b>B1FT</b> from <i>their</i> frequency table for heights of 0, 20, 80, 130, 210, 250 <b>B1FT</b> for increasing curve or polygon through 5 or 6 points
8(d)(ii)	1350 - 1400	2	<b>FT</b> <i>their</i> cumulative frequency diagram <b>M1</b> for correct UQ or LQ for <i>their</i> cumulative frequency diagram
9(a)	$3x^2 - 6x$	2	<b>M1</b> for $ax^2 - bx$ or <b>B1</b> for $3x^2$ or $-6x$
9(b)(i)	(0, 0) (2, -4)	4	M1 for their $ax^2 - bx = 0$ M1 for $x = 0$ and $x = \frac{theirb}{theira}$ or $x(ax-b) = 0$ A1 for $x = 0$ and $x = 2$ or $(0, 0)$ dep on first M1 seen or $(2, -4)$ dep on first M1 seen



Question	Answer	Marks	Partial Marks
9(b)(ii)	(0, 0) maximum (2, -4) minimum with supporting evidence	3	M1FT for $\left[\frac{d^2 y}{dx^2}\right] = 6x - 6$ or their $2ax - b$ A1 for $x = 0$ $\left[\frac{d^2 y}{dx^2}\right] = -6 < 0$ therefore maximum A1 for $x = 2$ $\left[\frac{d^2 y}{dx^2}\right] = 6 > 0$ therefore minimum Alternative method M1FT for considering $\frac{dy}{dx}$ both sides of x = 0 or $x = 2A1 for x = 0 is a maximum with validpoints tested correctlyA1 for x = 2 is a minimum with validpoints tested correctly$
9(c)	1 ±√3	3	M1FT for $3x^2 - 6x = 6$ or <i>their</i> $\frac{dy}{dx} = 6$ , dep on <i>their</i> $\frac{dy}{dx}$ being a function of $x$ . M1FT for $\frac{6 + \sqrt{(-6)^2 - 4 \times 3 \times (-6)}}{2 \times 3}$ or $\frac{6 - \sqrt{(-6)^2 - 4 \times 3 \times (-6)}}{2 \times 3}$ or better, dep on <i>their</i> $\frac{dy}{dx}$ being quadratic

