

Centre Number	Candidate Number	Name
---------------	------------------	------

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

PHYSICAL SCIENCE

0652/03

Paper 3

October/November 2005

1 hour 15 minutes

Candidates answer on the Question Paper.
No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen in the spaces provided on the Question Paper.
You may use a pencil for any diagrams, graphs, tables or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.
The number of marks is given in brackets [] at the end of each question or part question.
A copy of the Periodic Table is printed on page 16.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
Total	

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

This document consists of **14** printed pages and **2** blank pages.

1 Fig. 1.1 shows the arrangement of electrons in a lithium atom.




Fig. 1.1

(a) Lithium and potassium are both Group I metals. Complete the diagram in Fig. 1.2 to show the arrangement of electrons in a potassium atom.




Fig. 1.2

[2]

(b) When a small piece of lithium is dropped into a trough half filled with water a reaction takes place. Bubbles of the gas hydrogen are given off slowly and lithium hydroxide is formed.

(i) Write a balanced equation for this reaction.

..... [2]

(ii) Describe how you could prove that the gas given off is hydrogen.

test

.....

result

..... [2]

(c) A small piece of potassium is dropped into a trough half filled with water. Describe two differences that you would see between the reaction of lithium with water and that of potassium with water.

- 1.
.....
 - 2.
.....
- [2]

2 A ray of light enters a rectangular glass block at an angle of incidence of 66° . The glass has a refractive index of 1.45.

(a) Calculate the angle of refraction for this ray of light.
Write down the equation that you use and show all your working.

[3]

(b) Draw a fully labelled diagram to show the refraction of the light as it enters and leaves the glass block.

[3]

4

3 Copper(II) oxide reacts with dilute sulphuric acid.



In the preparation of copper(II) sulphate, copper(II) oxide is added to 20 cm³ of sulphuric acid of 1.0 mol/dm³ concentration until no more reacts.

(a) (i) Calculate the number of moles in the 20 cm³ of sulphuric acid.

moles of sulphuric acid = [1]

(ii) How many moles of copper(II) sulphate are produced in the reaction?

moles of copper(II) sulphate = [1]

(iii) Calculate the relative formula mass, *M_r*, of copper(II) sulphate, CuSO₄.

Show your working.

M_r = [2]

(iv) Calculate the mass of copper(II) sulphate, CuSO₄, formed.

Show your working.

mass =g [2]

(b) Describe how crystals of copper(II) sulphate can be prepared from the mixture of excess copper(II) oxide and copper(II) sulphate solution obtained when the reaction stops.

.....
.....
.....
..... [3]

- 4 A player throws a ball, of mass 0.15 kg, horizontally. The ball has a constant acceleration for a time of 0.10s and then moves at a constant speed of 20.0 m/s for 0.80 s before being caught and brought to rest in a further time of 0.30 s. As the ball is caught it decelerates non-uniformly.

- (a) On Fig. 4.1 draw a graph showing the speed of the ball from when it was thrown until the time it came to rest.




Fig. 4.1

[4]

- (b) Calculate the maximum kinetic energy of the ball. Show all your working.

maximum kinetic energy = [3]

- (c) Calculate the acceleration of the ball during the first 0.10 s. Write down the equation that you use and show all your working.

acceleration = [3]

5 Fig. 5.1 shows the gas hydrogen being burned in air.




Fig. 5.1

(a) When hydrogen burns the only product is water.
Write a balanced equation for the burning of hydrogen.

..... [2]

(b) When petrol is burned in a car engine a number of products are formed.
Some of these products cause pollution.
These include carbon monoxide and oxides of nitrogen.

(i) How are the oxides of nitrogen removed from the exhaust gases of modern cars.

..... [1]

(ii) Why may the presence of carbon monoxide in car exhaust systems cause a health problem?

..... [1]

(c) It has been suggested that hydrogen may replace petrol as a fuel for cars.
Suggest one advantage and one disadvantage of using hydrogen instead of petrol.

advantage

.....

disadvantage

..... [2]

6 (a) Explain what is meant by an object being in *equilibrium*.

.....
.....
..... [2]

(b) Fig. 6.1 shows a method of measuring the mass of a uniform loaded ruler. The ruler is pivoted at the 18 cm mark.




Fig. 6.1

(i) The ruler is uniform. What does this tell you about the position of its centre of mass?

.....
..... [1]

(ii) The total length of the ruler is 80 cm. The 50 g mass is hung from the 8 cm mark on the ruler. Calculate the mass of the ruler. Show all your working.

mass of ruler = g [4]

7 Powdered calcium carbonate is added to excess hydrochloric acid of three different concentrations, **A**, **B** and **C**.



In each experiment the same mass of powder is used and the acid is at the same temperature.

The volume of carbon dioxide gas given off is measured at time intervals.

The results of these experiments are shown in Fig. 7.1.




Fig. 7.1

(a) (i) Which of the three solutions of hydrochloric acid, **A**, **B** or **C**, is the most concentrated?

..... [1]

(ii) Explain how Fig. 7.1 shows your answer to (i) is correct.

.....
.....
..... [2]

(iii) Why do each of the three experiments give the same total volume of gas?

.....
..... [1]

(b) A fourth experiment is carried out using hydrochloric acid solution **A** and the same mass of powdered calcium carbonate.

This time the experiment is carried out at a higher temperature.

Sketch on Fig. 7.1 the result you would expect for this fourth experiment. [2]

(c) (i) Calculate the number of moles in the 100 cm³ of carbon dioxide gas produced.
(Assume the volume of carbon dioxide is measured at r.t.p. The volume of one mole of any gas is 24 dm³ at r.t.p.)

moles of carbon dioxide = [1]

(ii) Calculate the number of moles of calcium carbonate used to produce 100 cm³ of carbon dioxide gas.

moles of calcium carbonate = [1]

(iii) Calculate the mass of calcium carbonate used to produce 100 cm³ of carbon dioxide gas.
Show your working.
(The relative formula mass, M_r , of calcium carbonate = 100.)

mass of calcium carbonate = g [2]

8 (a) (i) Name the process by which the Sun produces energy.

.....


(ii) Explain what happens in this process.

.....
.....
.....
.....
.....
.....
..... [3]


(b) Calculate the energy released in the Sun when its mass decreases by 1200 kg as a result of this process. Write down the equation you use and show all your working. The speed of light = 3.0×10^8 m/s.

energy released = J [4]


9 Fig. 9.1 shows the graphical formulae of five organic compounds.




A




B



C



D



E

Fig. 9.1

(a) (i) Which **two** compounds are alkanes?

.....

(ii) Which compound dissolves in water to give an acidic solution?

.....

[1]

(b) (i) Describe a test to distinguish between compounds **C** and **D**.

test

.....

result

.....

[2]

(ii) In industry compound **D** is made from compound **C**.
Name the type of reaction that is used.

.....

[1]

(c) Compound **D** can be used to make a polymer.
Draw the structure for this polymer.

[2]

10 Fig. 10.1 shows a circuit with a high resistance voltmeter being used to measure the e.m.f. of a cell.




Fig. 10.1

(a) Explain why the voltmeter must have a high resistance if it is to measure an accurate value of the e.m.f.

.....
.....
..... [2]

(b) Fig. 10.2 shows a cell with an internal resistance of $5\ \Omega$. A voltmeter which has a resistance of $995\ \Omega$ is connected across the cell. The e.m.f. of the cell is 1.50 V .




Fig. 10.2

(i) Calculate the current in the circuit.

current = A [3]

13

(ii) Calculate the potential difference across the voltmeter.

potential difference = V [2]

(iii) Explain why this voltmeter gives a good approximation to the e.m.f. of the cell.

.....
.....
.....
..... [2]

DATA SHEET
The Periodic Table of the Elements

		Group															
I	II	III	IV	V	VI	VII	O										
1 H Hydrogen											2 He Helium						
3 Li Lithium	4 Be Beryllium	5 B Boron	6 C Carbon	7 N Nitrogen	8 O Oxygen	9 F Fluorine	10 Ne Neon										
11 Na Sodium	12 Mg Magnesium	13 Al Aluminium	14 Si Silicon	15 P Phosphorus	16 S Sulphur	17 Cl Chlorine	18 Ar Argon										
19 K Potassium	20 Ca Calcium	21 Sc Scandium	22 Ti Titanium	23 V Vanadium	24 Cr Chromium	25 Mn Manganese	26 Fe Iron	27 Co Cobalt	28 Ni Nickel	29 Cu Copper	30 Zn Zinc	31 Ga Gallium	32 Ge Germanium	33 As Arsenic	34 Se Selenium	35 Br Bromine	36 Kr Krypton
37 Rb Rubidium	38 Sr Strontium	39 Y Yttrium	40 Zr Zirconium	41 Nb Niobium	42 Mo Molybdenum	43 Tc Technetium	44 Ru Ruthenium	45 Rh Rhodium	46 Pd Palladium	47 Ag Silver	48 Cd Cadmium	49 In Indium	50 Sn Tin	51 Sb Antimony	52 Te Tellurium	53 I Iodine	54 Xe Xenon
55 Cs Caesium	56 Ba Barium	57 La Lanthanum	72 Hf Hafnium	73 Ta Tantalum	74 W Tungsten	75 Re Rhenium	76 Os Osmium	77 Ir Iridium	78 Pt Platinum	79 Au Gold	80 Hg Mercury	81 Tl Thallium	82 Pb Lead	83 Bi Bismuth	84 Po Polonium	85 At Astatine	86 Rn Radon
87 Fr Francium	88 Ra Radium	89 Ac Actinium															

140 Ce Cerium	141 Pr Praseodymium	144 Nd Neodymium	150 Sm Samarium	152 Eu Europium	157 Gd Gadolinium	162 Dy Dysprosium	165 Ho Holmium	167 Er Erbium	169 Tm Thulium	173 Yb Ytterbium	175 Lu Lutetium
90 Th Thorium	91 Pa Protactinium	92 U Uranium	94 Pu Plutonium	95 Am Americium	96 Cm Curium	98 Cf Californium	99 Es Einsteinium	100 Fm Fermium	101 Md Mendelevium	102 No Nobelium	103 Lr Lawrencium

*58-71 Lanthanoid series
90-103 Actinoid series

Key

a	X	a = relative atomic mass
	X	X = atomic symbol
b		b = proton (atomic) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).