UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the October/November 2006 question paper

0652 PHYSICAL SCIENCE

0652/06

Paper 6 (Alternative to Practical), maximum raw mark 60

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

The grade thresholds for various grades are published in the report on the examination for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2006 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

twww.xtrapapers.com

Total 10 marks

ge 2			Mark Scheme	Syllabu
J			IGCSE - OCT/NOV 2006	0652
1	(a)	(i) (ii)	carbon dioxide or CO ₂ lime-water, goes cloudy/white (precipitate) if an incorrect gas is named in (a)(i) a correct test f must be given	Syllable 1990 aper 0652 For that gas
	(b)	(i) (ii)	35.8 g, 39.2 g (no tolerance)(if given in wrong orde 3.4g (ecf)	er 1 mark only) [2]. [1]
	(c)		bubbling stops OWTTE (reject, all the solid has disanswers based on use of an indicator)	ssolved : any [1]
	(d)	(i) (ii)	26.2 g, 1.7 g (no tolerance) (if written in the wrong spaces 1 mark only) 24.5 g (ecf)	[2] [1]
	(e)		24.5 / 10 = 2.45g calcium carbonate (ecf) (1)	[1]
				Total 10 marks
2	(a)		12 mm, 63 mm: 17 mm, 60 mm (+/- 1 mm)	[4]
	(b)	(i)	axes correct & labelled (1) all points plotted accurately (+/- 1 mm)(1) straight line drawn cutting the y-axis (1) (no marks for graph of y (or x) against reading nun	nber) [3]
	(ii)		72 mm +/- 2 mm (answer checked with candidate's (no marks for answers given in centimetres)	s graph) [1]
	(iii)		line correctly drawn (+/- 1 mm) and point M labelle	d (ecf) [1]
	(c)		the weight of the card is not acting at point N/N is not the centre of mass/one side of the card is heavier than the other OWT	TE [1]

Page 2

www.xtrapapers.com

Page 3		Mark Scheme		Syllabu	per
	l	GCSE - OCT/NOV 20	006	0652	App Del
3 (2)				`	TOPANO.
3 (a)		sodium carbonate	ammonium chloride	aqueous ammonia	ambridge.
	! al	1		V	
	acid	V			

` '	sodium carbonate	ammonium chloride	aqueous ammonia
acid	V		V
base		V	
salt	V		V

		4 (or 5) correct (2) 2 or 3 correct (1) 1 or 0 (0)	[2]
		(note that if more than 4 ticks are shown, the number of wrong answers is subtracted from the number of correct answers)	
(b)		solid B is an acid/solid C contains a metal/solid A is not an acid/solid B produces carbon dioxide/other suitable conclusion	[1]
(c)	(i) (ii)	(moist) (red) litmus (Universal Indicator)(paper) (1) is turned blue(purple)/other suitable test for ammonia (1) solid A is a base (alkali)	[2] [1]
(d)	(i) (ii)	neutralisation/ammonia is neutralised/exothermic/salt formation metal (zinc) hydroxide (zinc need not be named for the mark) (accept aluminium)(reject metals giving a coloured hydroxide)	[1] [1]
(e)	(i) (ii)	add barium chloride (nitrate) (and hydrochloric (nitric) acid) (the addition of acid need not be mentioned) white solid/white precipitate seen	[1] [1]

Total 10 marks

WWW	Axtra	pa	per	s.cc	m
47.			•		

[1]

Total 10 marks

				2	
Page 4			Mark Scheme	Syllabl	per
			IGCSE - OCT/NOV 2006	0652	2
4	(a)		when the syringe reads a measured volume (100c clock (both statements are required for the mark)	m³), start the	Cannbridge Cop
	(b)		18, 43 cm ³ (no tolerance)		.col
	(c)		$CH_4 = 16$, $Cl_2 = 71$ (both correct)		[1]
	(d)		suitable named metal + hydrochloric or sulphuric a /electrolysis of suitable named electrolyte (reject "r		[1]
	(e)	(i)	B	the a warmen the a	[1]
		(ii)	the lower the rmm the faster the flow or the higher slower the flow (1) names and/or figures from the t		[2]
	(f)		no naked flame (1) hydrogen (methane) is flamma (1) or ventilate laboratory/use fume cupboard/do not chlorine is poisonous (1) reason must match safety precaution do not award mark for use of laboratory clothing, gother precaution not directly connected with the expectation of the connected with the connected with the expectation of the connected with the expectation of the connected with	ot inhale chlorine (1	[2]
				Т	otal 10 marks
5	(a)		73, 97°C, (2) 67, 73 cm³ (2) no tolerance		[4]
	(b)		sensible scale chosen and axes labelled (1) points plotted (1) +/- 1°C and 1cm ³ best straight line drawn (1)		[3]
	(c)		when heated molecules have more energy (1) molecules collide (with walls) (bounce) (or with each greater force (reject "vibrate")(1) collisions (between molecules) occur more often (•	
			volume or pressure is raised (1)	any 2 points	[2]

gas changes state/becomes liquid/solid (on cooling)

(d)

www.xtrapapers.com

Page 5	Mark Scheme Syllab.	aper
	IGCSE - OCT/NOV 2006 0652	TOO
6 (a)	52.5, 48.8, 47.3, 50.0 (must say 50.0) no tolerance	o aper aper apparation of the contract of the
(b)	B and C (1), D, A (in correct order) (1)	36.C
(c)	C, B (in correct order) ecf from (b)	
(d)	hydrocarbon will burn in air with a yellow (smoky) flame alcohol burns with a blue flame alcohol reacts with (conc) sulphuric acid alcohol will form an ester	F4
	alcohol will react with sodium/other suitable suggestion (any 1)	[1
(e)	add silver nitrate (solution) (1), white precipitate (1) OR yellow (orange) (1) flame test (1)	[2

Total 10 marks