

Cambridge Assessment International Education Cambridge International General Certificate of Secondary Education

PHYSICS

0625/42 March 2018

Paper 4 Extended Theory MARK SCHEME Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the March 2018 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is a registered trademark.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a guestion. Each guestion paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question •
- the specific skills defined in the mark scheme or in the generic level descriptors for the question •
- the standard of response required by a candidate as exemplified by the standardisation scripts. •

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded positively:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the • scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do •
- marks are not deducted for errors •
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the . question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

| March | 2018 |
|-------|------|
|-------|------|

| Question | Answer | Marks |
|----------|--|-------|
| 1(a) | Rate of change of velocity OR change of velocity / time OR change of velocity over time OR (v $-$ u)/t | B1 |
| 1(b)(i) | Straight line from origin to (15, 28) | B1 |
| | Horizontal line {from (15, 28)} to (32, 28) | B1 |
| | a = (v - u)/t OR (t =) (v - u)/a OR (0 - 28)/2.0 | C1 |
| | = 14 (s) | C1 |
| | Straight line from (32, 28) to (46, 0) | A1 |
| 1(b)(ii) | 1 Towards the centre of the circle / inwards | B1 |
| | 2 Velocity is (continually) changing its direction | B1 |

| Question | Answer | Marks |
|----------|---|-------|
| 2(a) | (Because g.p.e. is) the work done by the force OR the force × the distance that the object rises OR mgh and height is <u>greater</u> | B1 |
| 2(b) | mgh OR $80 \times 65 \times 10 \times 1600$ | C1 |
| | $8.3 \times 10^7 \text{ J}$ | A1 |
| 2(c) | Method 1 | ŀ |
| | W = Pt OR E = Pt in any form | C1 |
| | Work input =1500 \times 10 ³ \times 30 \times 60 OR 2.7 \times 10 ⁹ J | C1 |
| | Efficiency = work output / work input (x 100) | C1 |
| | 0.031 OR 3.1 % | A1 |
| | Method 2 | |
| | P = E/t in any form | (C1) |
| | Power output = $8.3 \times 10^7 / 30 \times 60$ | (C1) |
| | Efficiency = power output / power input (× 100) | (C1) |
| | 0.031 OR 3.1% | (A1) |

| Question | Answer | Marks |
|-----------|--|-------|
| 3(a) | (the) force × its perpendicular distance from pivot / a point | B1 |
| 3(b)(i) | centre of mass | B1 |
| 3(b)(ii) | (mass =) 160 / 10 = 16 kg | B1 |
| 3(b)(iii) | (Not moving up or down because) no resultant (vertical) force OR upward force = downward force | C1 |
| | 80 N + 80 N = 160 N | A1 |
| | (Not rotating because) no resultant moment (about any point) OR (sum of) clockwise moments = (sum of) anticlockwise moments | C1 |
| | clockwise moment (about centre) = 80×1.2 anticlockwise moment (about centre) = 80×1.2 | A1 |

| Question | Answer | Marks |
|----------|--|-------|
| 4(a) | Molecules of hot liquid collide with (surface of) spoon | B1 |
| | transfer energy / heat to (molecules of) spoon | B1 |
| | (amplitude of) vibration of spoon's molecules increases / is faster (increasing spoon's temperature) | B1 |
| 4(b) | Molecules of hot liquid (also) transfer energy to (free) electrons in the spoon | B1 |
| | These (free) electrons move through the metal | B1 |
| 4(c) | $(Q =) mc\Delta\theta$ | C1 |
| | 150 × 4.2 × (80 – 56) | C1 |
| | 15000 J | A1 |

| Question | Answer | Marks |
|----------|--|-------|
| 5(a)(i) | Sketch showing straight lines with sudden changes of direction | B1 |
| 5(a)(ii) | Any 3 marks from 4 points: | |
| | Air molecules move in random / different directions | B1 |
| | Smoke particles are hit | B1 |
| | by air molecules | B1 |
| | Change direction at each collision OR undergo Brownian motion | B1 |
| 5(b) | F = (mv – mu) / t in any form OR Impulse = mv – mu | C1 |
| | $= 20 \times 4.2 / 60$ | C1 |
| | 1.4 N | A1 |

| Question | Answer | Marks |
|-----------|--|-------|
| 6(a) | A: infra-red B: ultra-violet C: X-(rays) D: γ-(rays) | B2 |
| 6(b)(i) | n = sin i / sin r OR sin r = sin i / n OR sin r = sin 35 / 1.50 | C1 |
| | r = 22° | A1 |
| 6(b)(ii) | Refraction at XY drawn with r < i | B1 |
| | Refraction at XZ drawn with r > i | B1 |
| 6(b)(iii) | Blue ray drawn below red ray in prism and drawn with r < i | M1 |
| | Ray to right of prism diverging downwards from red ray | A1 |

| Question | Answer | Marks |
|-----------|---|-------|
| 7(a) | n = speed in air / speed in water OR speed in water = 3.0×10^8 / 1.33 | C1 |
| | $2.3 \times 10^8 \text{m/s}$ | A1 |
| | Answers to (b)(i), (b)(ii) and (b)(iii) all combined to maximum of 5 marks on same screen | |
| 7(b)(i) | Wavefronts in plastic: meet wavefronts in air | B1 |
| | make smaller angle with boundary than wavefronts in air and downwards to the left | B1 |
| | parallel to each other | B1 |
| 7(b)(ii) | Arrow(s) perpendicular to wavefronts in plastic and downwards to right | B1 |
| 7(b)(iii) | r in plastic between refracted wavefront and boundary | B1 |
| | OR At a point where refracted wavefront meets boundary, normal to boundary drawn and line perpendicular to wavefront drawn. <i>r</i> in plastic between the two lines drawn | (B1) |

www.xtrapapers.com March 2018

| Question | Answer | Marks |
|----------|--|-------|
| 8(a) | P = IV OR (I =) 50 / 12 | C1 |
| | 4.2 A | A1 |
| 8(b)(i) | (E =) QV | C1 |
| | $(E =) 270 \times 10^3 \times 12$ | C1 |
| | $3.2 \times 10^{6} \text{J} / 3200 \text{kJ}$ | A1 |
| 8(b)(ii) | Volume of fuel used = $3.2 \times 10^6 / 3.6 \times 10^4$ | C1 |
| | 89 cm^3 OR 90 cm ³ if 3.24×10^6 used | A1 |

| Question | Answer | Marks |
|----------|--|-------|
| 9(a)(i) | Resistor: tick in 2nd box | B1 |
| 9(a)(ii) | Lamp: tick in 1st box | B1 |
| 9(b) | (R =) V/I OR (R =) 6.0/4.4 | C1 |
| | 1.4 Ω | A1 |
| 9(c) | Current in lamp = 4.4 A Current in resistor = 4.0 A | C1 |
| | Current from supply (= 4.0 + 4.4) = 8.4 A | A1 |
| | OR (With 6 V p.d.) $R_L = 6/4.4 = 1.36 \Omega R_R = 6/4 = 1.5 \Omega$ Combined resistance = $(1.36 \times 1.5)/2.86$) = 0.71 Ω | (C1) |
| | Current = 6 / 0.71 = 8.4 A | (A1) |
| 9(d) | p.d. across lamp = 4.9 V p.d. across resistor = 6.0 V | C1 |
| | Total p.d. (= 4.9 + 6.0) = 10.9 V | A1 |
| | OR (With 4 A current) $R_L = 5/4 = 1.25 \Omega R_R = 6/4 = 1.5 \Omega$ Total R = 2.75 Ω | (C1) |
| | Total p.d. = 2.75 × 4 = 11.0 V | (A1) |

| Question | Answer | Marks |
|-----------|---|-------|
| 10(a) | Strength of magnetic field / magnet | B1 |
| | Speed (of movement of wire) | B1 |
| | Length of AB / wire (within field) | B1 |
| 10(b)(i) | $V_p / V_s = N_p / N_s$ OR (N _s =) 8000 × 12 / 240 | C1 |
| | 400 (turns) | A1 |
| 10(b)(ii) | Circuit connected to A and B with resistor and diode with correct circuit symbols in series | B1 |

| Question | Answer | Marks |
|-----------|---|-------|
| 11(a)(i) | In box / cupboard with lead walls | B1 |
| 11(a)(ii) | (Handle) with (long) tongs OR remote-controlled device OR wearing lead gloves OR wearing lead suit | B1 |
| 11(b) | Col 1: gamma / γ (rays) alpha / α (particles) beta / β (particles) | B1 |
| | Col 3: – a few cm or up to 10 cm a few m or up to 10 m | B1 |
| | Col 4: thick lead or 30 cm lead or very thick concrete or 3 m concrete - thin aluminium or 2 mm aluminium | B1 |
| 11(c)(i) | alpha / α (particles or rays) | B1 |
| 11(c)(ii) | beta / β (particles or rays) | B1 |