

Cambridge IGCSE™

PHYSICS

Paper 3 Core Theory MARK SCHEME Maximum Mark: 80 0625/33 May/June 2022

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2022 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question •
- the specific skills defined in the mark scheme or in the generic level descriptors for the question .
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the • scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do .
- marks are not deducted for errors •
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the • question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.

5 <u>'List rule' guidance</u>

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards **n**.
- Incorrect responses should not be awarded credit but will still count towards *n*.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 <u>Calculation specific guidance</u>

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 <u>Guidance for chemical equations</u>

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Α	1	Correct	~	2	F	1	Correct	✓	2
	2	Correct	✓		(4 responses)	2	Correct	√	
	3	Wrong	×			3	Correct CON (of 3.)	× (discount 3)	
В	1	Correct, Correct	√, √	3					
(4 responses)	2	Correct	✓		G	1	Correct	✓	3
(3		ianoro		(5 responses)	2	Correct	✓	
	<u></u> з	Wrong	ignore			3	Correct Correct	√ ignore	
С	1	Correct	✓	2			CON (of 4.)	ignore	
(4 responses)	2	Correct, Wrong	√, ×						
	3	Correct	ignore		н	1	Correct	✓	2
					(4 responses)	2	Correct	*	
D	1	Correct	✓	2		3	CON (of 2.) Correct	(discount 2) ✓	
(4 responses)	2	Correct, CON (of 2.)	×, (discount 2)						
	3	Correct	\checkmark		1	1	Correct	✓	2
				1	(4 responses)	2	Correct	×	
E	1	Correct	✓	3		3	Correct	√	
(4 responses)	2	Correct	✓				CON (of 2.)	(discount 2)	
	3	Correct, Wrong	✓						

M marksare method marks upon which further marks depend. For an M mark to be scored, the point to which it refers **must** be seen in a candidate's answer. If a candidate fails to score a particular M mark, then none of the dependent marks can be scored.

- B marks are independent marks, which do not depend on other marks. For a B mark to scored, the point to which it refers must be seen specifically in the candidate's answers.
- A marks In general A marks are awarded for final answers to numerical questions. If a final numerical answer, eligible for A marks, is correct, with the correct unit and an acceptable number of significant figures, all the marks for that question are normally awarded. It is very occasionally possible to arrive at a correct answer by an entirely wrong approach. In these rare circumstances, do not award the A marks, but award C marks on their merits. However, correct numerical answers with no working shown gain all the marks available.
- C marks are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate, **provided subsequent working gives evidence that they must have known it.** For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows that they knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.
- Brackets () around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.
- <u>Underlining</u> indicates that this <u>must</u> be seen in the answer offered, or something very similar.
- OR / or indicates alternative answers, any one of which is satisfactory for scoring the marks.
- o.w.t.t.e. means 'or words to that effect'.

Question	Answer	Marks
1(a)(i)	metre rule	B1
1(a)(ii)	50 (m)	B1
1(b)	graph starts at origin	B1
	speed = 0 at 9.0 s	B1
	highest speed at 17 s	B1
1(c)(i)	260 (s)	B1
1(c)(ii)	1.9 (m/s)	A3
	500 ÷ 260 OR 500 ÷ (c)(i)	(C2)
	(speed =) distance ÷ time in any form	(C1)

Question	Answer	Marks
2(a)(i)	electrical	B1
	(to) kinetic OR (gravitational) potential	B1
2(a)(ii)	(gravitational) potential	B1
2(b)(i)	same	B1
2(b)(ii)	any two from the following: energy transferred to surroundings thermal (energy) / heat lost work done against friction motor less than 100% efficient	B2

Question	Answer	Marks
2(b)(iii)	weight/force/mass AND height/distance moved	B1
	time	B1
2(c)	7900 OR 7870	A3
	70 ÷ 0.0089	(C2)
	(density =) mass ÷ volume in any form	(C1)

Question	Answer	Marks
3(a)	any two from larger area lower pressure (on ground) does not sink in	B2
3(b)(i)	400	A3
	4000 ÷ 10 OR 4000 ÷ 9.8	(C2)
	(mass =) weight ÷ g OR weight ÷ 10 weight ÷ 9.8 in any form	(C1)
3(b)(ii)	500	A3
	1000 ÷ 2.0 OR 4000 ÷ (4 × 2.0)	(C2)
	(pressure =) force ÷ area in any form	(C1)
	N/m ² OR Pa	B1

Question	Answer	Marks
4(a)	situation	B1
	explanation	B1
4(b)(i)	decreases	B1
4(b)(ii)	molecules slow(er)/less (kinetic) energy	B1
	hit (wall/flask) <u>less</u> often	B1
	hit (wall/flask) with <u>less</u> force	B1

Question	Answer	Marks
5(a)	thermometer	B1
5(b)(i)	radiation	B1
5(b)(ii)	conduction	B1
5(c)	(dull) black pipes	B1
	better absorber (of radiation)/less reflection from pipes	B1
	foil/shiny metal behind pipes	B1
	reflects (radiation back to pipes) owtte	B1
	valid alternatives	(B1)

Question	Answer	Marks
6(a)	normal(s)	B1
6(b)(i)	angle of incidence	B1
	that gives an angle of refraction of 90°	B1
6(b)(ii)	20° – emergent ray is refracted in air	B1
	40° – emergent ray is refracted in air AND greater refraction	B1
	60° – ray is reflected internally	B1

Question	Answer	Marks
7(a)(i)	infrared AND microwaves AND radio waves	B1
7(a)(ii)	radio waves	B1
7(a)(iii)	any two from: infrared microwaves radio waves	B1
7(a)(iv)	ultrasound	B1
7(b)	remote controls/heat lamps/heaters/thermal imaging/night vision/security cameras/heaters/grills/measuring blood	B1
	oxygen/pulse oximeters/measuring body temperature/thermal thermometer	B1

Question		Answer		Marks
7(c)	wave	transverse	longitudinal	B2
	infrared	✓		
	microwaves	✓		
	ultrasound		✓	
	radio waves	×		
		1	1	

Question	Answer	Marks
8(a)(i)	400	A3
	$(Ns =) N_P \times V_S / V_P OR 8000 \times 12/240$	(C2)
	$V_P/V_S = N_P/N_S$ in any form	(C1)
8(a)(ii)	(step-down) transformer	B1
8(a)(iii)	copper	B1
8(b)	motor connected between A and B	B1
	2 lamps in series connected between A and B	A2
	2 lamps in series	(C1)

Question	Answer	Marks
9(a)	electromotive force/e.m.f./ potential difference/p.d./ voltage	B1

Question	Answer	Marks
9(b)(i)	electron(s)	B1
9(b)(ii)	ammeter	B1
9(c)(i)	0.4(0) (A)	A3
	4(.0) ÷ 10	(C2)
	$(I =) \vee \div R$ in any form	(C1)
9(c)(ii)	0.4(0) (A) OR same as (c)(i)	B1
9(c)(iii)	no change/same	M1
	resultant/total/combined resistance does not change	A1

Question	Answer	Marks
10(a)	alpha/α	B1
	beta/β	B1
	gamma/γ	B1
10(b)(i)	35	B1
10(b)(ii)	17	B1
10(b)(iii)	18	B1
10(c)	electrons	B1