

**CANDIDATE** NAME

## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

 CENTRE NUMBER
COMBINE
Paper 6 A

**COMBINED SCIENCE** 

0653/61

Paper 6 Alternative to Practical

May/June 2010

CANDIDATE NUMBER

1 hour

Candidates answer on the Question paper

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs, tables or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [ ] at the end of each question or part question.


For Examiner's Use		
1		
2		
3		
4		
5		
6		
Total		

This document consists of 19 printed pages and 1 blank page.



- 1 One of the characteristics of living things is the ability to detect changes surroundings and respond to them. This is important for survival.
- WANN. P. BRAC CAMBridge. COM (a) A choice chamber was used to study the responses of woodlice (small invertebrates) to the intensity of light and the amount of moisture in their surroundings. The choice chamber is shown in Fig. 1.1.





- Fig. 1.1
- The divider separated the base of the chamber in half. One side was dry and the other side was damp.
- A platform for the woodlice was made from muslin (a type of net) and this covered the base.
- Half of the lid was covered with light-proof paper. The lid was placed so that four different conditions were produced. The conditions were:

dry and dark,

damp and dark,


dry and light,

damp and light.

- Ten woodlice were placed on the centre of the muslin and left for 15 minutes.
- After this time the number of woodlice in each area was counted.

The experiment was done three times. After each test the lid was removed and the number of woodlice in each area was counted. The platforms when viewed from above are shown in Fig. 1.2.

(i) Count the number of woodlice in each area of the choice chamber and en numbers in Table 1.3. Test A has been done for you.







Fig. 1.2

Table 1.3

conditions in	numbe	number of woodlice counted			
each area	test A	test <b>B</b>	test C	average number of woodlice	
dry and dark	2				
damp and dark	6				
damp and light	2				
dry and light	0				

(ii) Calculate the average numbers of woodlice in each area and complete the last column of Table 1.3. [2]





[3]

(c)	(i)	Which <b>two</b> external conditions did the woodlice prefer?	
		and	[1]
	(ii)	Choose <b>one</b> of these conditions and suggest how this may help the woodlice survive.	to
			 [2]

2 A student is investigating the acceleration of a falling mass due to the force of gravity using a ticker-timer. This apparatus is shown in Fig. 2.1.




Fig. 2.1

(a)

The iron bar is fixed at one end. The bar passes between the poles of a permanent magnet. An alternating current passing through the coil causes the bar to vibrate 50 times per second. At the other end of the bar the pin hits the carbon paper disc making a mark on the paper strip each time the bar vibrates.

(i)	Explain what is meant by an alternating current.	
		[1]
(ii)	Explain why the alternating current makes the iron bar vibrate.	
		[2]




Table 2.3

6		WWW.	xtrapapers.com  Aba  For iner's
	Table 2.3		For iner's
time/s	distance from point 0 / cm	distance from point 0 / m	Tidde com
0.00	0.0	0.000	
0.02	0.2	0.002	
0.04	0.7	0.007	
0.06	1.7	0.017	
0.08	3.1	0.031	
0.10	4.8	0.048	
0.12	6.9	0.069	
0.14			
0.16			
0.18			

Fig. 2.2

A 100 g mass is fixed on the end of the strip of paper. The student allows the mass to fall. It pulls the paper through the ticker-timer as shown in Fig. 2.1. Marks are made on the paper strip.

Part of the paper strip is shown in Fig. 2.2. The student is measuring the distance of each mark from the start, point 0. He has recorded the first six distances in Table 2.3.

- (b) (i) Using Fig. 2.2, measure the distances from point 0 to the dots at  $t = 0.14 \, \text{s}$ , 0.16 s, and 0.18 s, in centimetres, to the nearest millimetre. Record the distances in Table 2.3. [3]
  - (ii) Complete Table. 2.3 by converting these three distances from centimetres to metres. [1]

(	(iii) Use	e data from	Table 2.3	s to show tha	at the mas	ss and paper	accelerated	d as the	Cam
									No.
									[1]
(c)		te <b>g</b> , the acg	cceleratio	on due to g	ravity, by	using data	from Table	e 2.3 and	the
			g =	2 x distance (time of	<u>e of fall in</u> fall in sec				
						g =	m/s	s <sup>2</sup>	[2]

The science teacher gives a student the apparatus shown in Fig. 3.1. She also gives 3 three solutions, **X**, **Y** and **Z** of different concentrations of the same acid.

The student must find out which of the three acid solutions is the most concentrated.




Fig. 3.1

- The student places  $5\,\text{cm}^3$  of the acid  $\boldsymbol{X}$  in the beaker.
- She adds 1 drop of the indicator.
- She adds the alkali, 1 drop at a time, until the indicator changes colour.
- She records the number of drops of alkali added in Table 3.2.
- She repeats the experiment with the acid solutions Y and Z.

Table 3.2

acid solution	number of drops
x	22
Y	17
Z	8

Table 3.3 below shows the colour of the indicator at different levels of pH.


Table 3.3

colour of indicator	red	orange
pH of solution	5	9

(a)	State the colou	State the colour of the indicator		
	in acid,			
	in alkali.		[1]	

(b)	(i)	Which of the acid solutions, <b>X</b> , <b>Y</b> or <b>Z</b> is the most concentrated?	aCambrida and an
			Ti
	(ii)	Explain your answer.	
			[1]
(a)	Suc	agost why the heaker of water is needed	
(C)	Sug	ggest why the beaker of water is needed.	
	•••••		[1]
(d)	Nar	me an alkali that can be used in this experiment.	
			[1]
2 cı	e stu m³ o ned.	dent wants to find out the name of the acid used in the experiment. She plant f solution <b>X</b> in a test-tube and adds aqueous silver nitrate. A white precipita	aces te is
(e)	(i)	Name the white precipitate.	
			[1]
	(ii)	Name the acid in solution <b>X</b> .	
			[1]
(f)		scribe how you could use pieces of magnesium ribbon to find out which of the tutions is the most concentrated.	hree
			[3]

A student did an experiment to find the density of salt solution. He floated a tecontaining sand, in water and then in the salt solution. This is shown in Fig. 4.1.







Fig. 4.1


- He placed dry sand in the test-tube and floated it in water. He measured the depth  $d_1$  from the water surface to the bottom of the tube. He recorded this in the first row of Table 4.2, **experiment 1**.
- He placed the same tube in the salt solution and found the depth  $d_2$ . He recorded this in the first row of Table 4.2.
- He emptied some of the sand out of the test-tube and found  $d_1$  and  $d_2$  again, experiment 2.
- He emptied out more of the sand and found another set of readings for  $d_1$  and  $d_2$ , experiment 3.

Table 4.2

experiment number	depth <b>d</b> ₁ in water/ millimetres	depth <b>d₂</b> in salt solution/millimetres
1	104	86
2	85	70
3		

Fig. 4.3 shows the tube floating in water and in salt solution for experiment 3. The has placed a ruler graduated in millimetres next to the tube.





tube floating in salt solution

Fig. 4.3

(a)	(1)	beaker.	ıne
			 [1]
(ii)		Use the scale of the ruler to calculate the depth $d_1$ for the tube floating in water.	
		Record the value in Table 4.2.	


depth  $d_1$  = \_\_\_\_\_mmm

(iii) Use the scale of the ruler to calculate the depth  $\mathbf{d_2}$  for the tube floating in salt solution.

Record the value in Table 4.2.

depth 
$$d_2 =$$
 mm [1]

(b) On the grid below, plot the values for  $d_1$  from Table 4.2 (vertical axis) against  $d_2$ the best straight line and extend it to pass through the point (0,0).



[2]

(c) Calculate the gradient of the line, showing on your graph the values you use to do this. The gradient is numerically equal to the density of the salt solution in grams per cubic centimetre.

> gradient = [2]

(d)	Describe another method for finding the density of a liquid using a pipette or bubbeaker and a balance.	For iner's
		G. COM
		`
	[2]	

5 Some plants show differences between leaves found in sunny areas of the plants leaves), and leaves found in shaded areas (shade leaves).

(a) A student was supplied with two leaves, labelled sun leaf and shade leaf. Drawings of the two leaves are shown in Fig. 5.1.

(i) Measure the maximum length of each leaf, excluding the petiole (stalk). Write your measurements below each diagram.

(ii)






Fig. 5.1

	Suil leal	Silade leal	
	length = mm	length = mm	[2]
(ii)	The leaves clearly have different areas.		
	One leaf has a larger surface than the other.		
	Suggest an advantage to the leaf with the la	ger area.	
			[1]

[4]

(b) The diagrams in Fig. 5.2 show cross-sections of a sun leaf and a shade leaf as using a microscope.




Fig. 5.2

Construct a table to compare these two diagrams. Include the following features; thickness of leaf, number of palisade cells, size of air spaces.

(c)	Choose any <b>one</b> feature of the sun leaf in Fig. 5.2 and explain how this feature is good adaptation for photosynthesis.
	feature
	explanation

(d)	The sun leaf usually has a thicker cuticle than the shade leaf. The cuticle is layer covering the leaf.	For iner's
	Suggest an advantage that this thicker cuticle gives to the sun leaf.	Tage
		[1]

6 The science teacher is showing the class an experiment. He makes carbon dioxic converts it into carbon monoxide. Then he reacts the carbon monoxide copper(II) oxide.

Fig. 6.1 shows the apparatus.




Fig. 6.1

(a)	Name a solid and a liquid that can be placed in the flask to make carbon dioxide.			
	solid			
	liquid			[2]
(b)	Comp	plete the symbol equation to show the reaction in the s	steel tube.	
		+	→ 2CO	[1]

WWW. Papa Cambridge Com (c) When the copper(II) oxide in the glass tube reacts with carbon monoxide it c from black to reddish-brown. The teacher takes the dish out of the glass tube who has cooled. He uses the apparatus shown in Fig. 6.2 to find out if the reddish-brown residue in the dish is a metal.




Fig. 6.2

Suggest two observations that the class can make, which show that the residue is a metal.

١	
2	[2]

(d) Before the experiment began, the teacher weighed the empty dish. Then he weighed it containing the copper(II) oxide. He recorded the masses in Table 6.3.

Table 6.3

mass of empty dish/g	mass of dish + copper(II) oxide/g	mass of dish + reddish-brown residue/g
35.9	43.9	

He weighs the dish containing the reddish-brown residue. Fig. 6.4 shows the balance window.




Fig. 6.4

(i) Read the balance window. Record the mass of the dish containing the reddish-brown residue in Table 6.3. [1]

	(ii)	Use the data from Table 6.3 to calculate the mass of the copper(II) oxide teacher placed in the dish.
		mass of copper(II) oxide = g [1]
	(iii)	Calculate the loss in mass of the copper(II) oxide when it reacted with carbon monoxide.
		loss in mass = g [1]
	The	class decide that oxygen has been removed from the copper(II) oxide.
	(iv)	What name is given to this process?
		[1]
(e)		ggest a reason why the teacher carried out this experiment in a fume cupboard ead of on the open laboratory bench.
		[1]

20

**BLANK PAGE** 

\*\*Www.xtrapapers.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.