	Cambridge IGCSE	Cambridge International Examinations Cambridge International General Certificate of S	Secondary Educ	ation
	CANDIDATE NAME			
* 3 6	CENTRE NUMBER		CANDIDATE NUMBER	
	COMBINED SC	IENCE		0653/51
7 2	Paper 5 Practic	al Test	Oc	tober/November 2015
6918				1 hour 30 minutes
	Candidates ans	wer on the Question Paper.		
3 1 7 J	Additional Mate	rials: As listed in the Confidential Instructions.		
* 💻	READ THESE	INSTRUCTIONS FIRST		

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units. Notes for Use in Qualitative Analysis for this paper are printed on page 8.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
Total		

This document consists of 8 printed pages.

CAMBRIDGE International Examinations


[Turn over

- **1** You are going to investigate what happens when food is burned in air.
 - (a) (i) Using the forceps, pick up a piece of dry cobalt chloride paper.
 Cobalt chloride paper changes colour from blue to pale pink or white in the presence of water.
 - Remove the stopper from one of the large test-tubes provided and place the piece of blue cobalt chloride paper inside.
 - Wait five seconds and then record the colour of the cobalt chloride paper.

[1]

- Remove the cobalt chloride paper from the test-tube.
- (ii) Add 10 cm³ limewater to the same test-tube used in part (i). Replace the stopper and shake gently for 10 seconds. Describe the appearance of the limewater.

(b) • Remove the stopper from the second test-tube and grip it with the test-tube holder as shown in Fig. 1.1.





- Place the piece of food provided on the mounted needle.
- Using a Bunsen burner, ignite the piece of food. Move the Bunsen burner to one side. Collect any gases produced by holding the second large test-tube inverted about 2 cm above the flame produced as shown in Fig. 1.1.
- Wait until the sides of the test-tube mist up and then place the food into the beaker of water to extinguish it.
- Replace the stopper on the test-tube.

© UCLES 2015

Place a fresh piece of blue cobalt chloride paper into the test-tube and quickly replace the stopper. Wait five seconds and then describe and explain the appearance of the cobalt chloride paper. appearance explanation[1] (c) Remove the piece of cobalt chloride paper and quickly add about 10 cm³ limewater to the test-tube. Replace the stopper and shake the test-tube for 10 seconds. (i) Describe the appearance of the limewater. [1] (ii) Explain your observation. [1] _____ (d) State two observations of this experiment that show energy is released. (e) State the purpose of the tests in part (a). [1] _____ (f) In this experiment you burned some food in air. Name the process inside living cells that this procedure is modelling. [1] _____ (g) State one precaution you took to make sure that the procedure in part (b) was carried out safely. [1]

[Turn over

www.xtrapapers.com

2 Potassium iodide can be converted to iodine by an oxidising agent.

The presence of starch produces a blue-black colour when the iodine is formed.

The time for the blue-black colour to appear depends on the rate of the reaction between potassium iodide and the oxidising agent.

You are going to investigate whether certain metal ions can catalyse this reaction.

A is potassium iodide solution.

B is a solution of the oxidising agent.

- (a) (i) Using the measuring cylinder labelled **B**, place 10 cm³ of **B** in a conical flask or beaker.
 - Add five drops of starch to the flask.
 - Using a second measuring cylinder labelled **A**, add 10 cm³ of **A** to the flask and swirl once, starting the stopclock at the same time.
 - Stop the stopclock when the solution turns blue-black. Record the time *t* in seconds to the nearest second in the second row of Table 2.1. [1]

volume of B / cm ³	drops of starch	volume of A /cm ³	1 cm ³ of solution of metal ion	time <i>t</i> /s
10.0	5	10.0	none added	
10.0	5	10.0	none added	
10.0	5	10.0	Fe ²⁺	
10.0	5	10.0	Fe ³⁺	

Table 2.1

- (ii) Repeat (a)(i) and record the time *t* in seconds to the nearest second in the next row of Table 2.1. [2]
- (b) (i) Using the measuring cylinder labelled **B**, place 10 cm³ of **B** in a conical flask or beaker.
 - Add five drops of starch to the flask.
 - Using the measuring cylinder labelled **C** add 1 cm³ of the solution containing the Fe²⁺ ion to the flask.
 - Using the measuring cylinder labelled **A**, add 10 cm³ of **A** to the flask and swirl once, starting the stopclock at the same time.
 - Stop the stopclock when the solution turns blue-black. Record the time *t* in seconds to the nearest second in the appropriate row of Table 2.1. [1]

© UCLES 2015

- 5
- (ii) Repeat (b)(i), replacing the solution containing the Fe^{2+} ion with the solution containing the Fe^{3+} ion.

Record the time *t* in seconds to the nearest second in the appropriate row of Table 2.1. [1]

(c) Using your results in Table 2.1, state and explain whether the metal ions are acting as catalysts for this reaction.

(d) Use the two results obtained when no metal ion was added to comment on the reliability of the experiment. You must show how you use these two results.

(e) The addition of the metal ion in (b) increases the total volume of the solution in the experiment.

Suggest a modification to the method in (a)(i) which would result in the overall experiment being a fairer test.

Explain why you have chosen this modification.

[2]

[Turn over

- **3** You are going to measure the capacity of a plastic cup.
 - (a) The capacity of a cup is the maximum volume of liquid that it can hold. The volume will be estimated by finding the average diameter of the cup and then considering the cup to be an approximate cylinder.
 - (i) Measure, to the nearest 0.1 cm, the height *h* of the cup as shown in Fig. 3.1.




Fig. 3.1

		h =	cm
(ii)	Measure the diameter <i>D</i> of the top of the cup.	D =	cm
(iii)	Measure the diameter <i>d</i> of the bottom of the cup.	d =	cm

(iv) Calculate the average diameter d_A using your results from (a)(ii) and (a)(iii) and the equation

$$d_{\mathsf{A}} = \frac{(D+d)}{2}$$

*d*_A = _____cm

(v) Calculate the approximate volume V of the cup using the equation

$$V = \frac{\pi d_A^2 h}{4}$$

V =_____ cm³ [6]

0653/51/O/N/15

© UCLES 2015

www.xtrapapers.com

(b)	(i)	Fill the measuring cylinder with water up to a mark in excess of 200 cm ³ .			
		Record this reading R ₁ .	R ₁ =	cm ³	
		Pour water from the measuring cylinder into the c	up until it is full.		
		Record the new reading R ₂ .	R ₂ =	_c cm ³	
		Determine the volume of water $V_{\rm W}$ that the cup ca	an hold. Show your working.		
			<i>V</i> _W =	cm ³	[2]
	(ii)	Suggest one possible source of inaccuracy in this	s procedure.		
				,	[1]
	(iii)	State which of the two values, V obtained in (a) accurate.	(v) or V _w obtained in (b)(i) i	is the m	ore
		Explain your answer.			
					[1]

NOTES FOR USE IN QUALITATIVE ANALYSIS

Test for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify then add aqueous barium chloride <i>or</i> aqueous barium nitrate	white ppt.

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
ammonium (NH ₄ ⁺)	ammonia produced on warming	-
copper(II) (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Test for gases

gas	test and test results
ammonia (NH ₃)	turns damp red litmus paper blue
carbon dioxide (CO ₂)	turns limewater milky
chlorine (Cl ₂)	bleaches damp litmus paper
hydrogen (H ₂)	"pops" with a lighted splint
oxygen (O ₂)	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.