



Cambridge International Examinations
Cambridge Pre-U Certificate

FURTHER MATHEMATICS

9795/02

Paper 2 Further Applications of Mathematics

May/June 2016

MARK SCHEME

Maximum Mark: 120

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 3 Pre-U Certificate.


This document consists of **7** printed pages.

Page 1	Mark Scheme	Syllabus	Paper		
	Cambridge Pre-U – May/June 2016	9795	02		
1	(i)	$75 \pm 1.96 \sqrt{\frac{40^2}{500} \times \frac{500}{499}}$ $= (71.5, 78.5)$	M1 B1 A1 A1 [4]	$75 \pm zs$, s involving 500 $z = 1.96$, allow from no 500 Variance correct Both limits correct to 3sf	<div style="border: 1px solid black; padding: 5px; width: fit-content;"> Condone omission of $\frac{500}{499}$ </div>
	(ii)	No, as the Central Limit Theorem applies OR as n is large	B1 [1]	“No” and mention CLT or large sample size; focus on different distributions; no irrelevancies	
2	(i)	$N(120, \dots)$ $\sigma^2 = 0.8^2 \times 1200 [= 768]$ $1 - \Phi((140 - 120)/\sqrt{768})$ $= 0.235(3)$	M1 M1 A1 A1 [4]	Normal, mean 120 or 1.20 Allow 0.8×1200 etc Both parameters correct Answer, in range [0.235, 0.236] OR: $P(\geq 175)$ from $N(150, 200)$ M1 $(175 - 150)/\sqrt{1200}$ A1 $0.235(3)$ A2	
	(ii)	$B_1 + B_2 + B_3 + B_4 - S_1 - S_2 - S_3$ $\sim N(60, \dots)$ Variance $4 \times 1200 + 3 \times 1500 = 9300$ $\Phi\left(\frac{0 - 60}{\sqrt{9300}}\right) = \Phi(-0.622) = 0.267$	M1 M1 A1 A1 [4]	Consider $\pm(B_1 + B_2 + B_3 + B_4 - S_1 - S_2 - S_3)$ or $4B - 3S$ Normal, mean 60 Correct variance Answer, a.r.t. 0.267 [0.2699] [NB: $\Phi(-60/\sqrt{33700}) = 0.3700$ is 2/4]	
3	(i)	$\sum t^r P(r) = \sum_{r=0}^n t^r {}^n C_r p^r (1-p)^{n-r}$ $= \sum_{r=0}^n (pt)^r {}^n C_r (1-p)^{n-r}$ $= (1-p + pt)^n \quad \text{AG}$	M1 A1 A1 [3]	Use $\sum t^r P(R=r)$ and binomial probabilities Indicate correct final term Collect p^r and t^r and correctly obtain given expression OR $(1-p + pt)^1$, M1A1; answer, A1	
	(ii)	$\left(\frac{3}{4} + \frac{1}{4}t\right)^8 \left(\frac{1}{4} + \frac{3}{4}t\right)^8$ $= (3+t)^8 (1+3t)^8 / 4^{16}$ $= \left(\frac{3}{16} + \frac{1}{16}t(10+3t)\right)^8 \quad \text{AG}$ $= \left(\frac{3}{16}\right)^8 \left(1 + \left[\frac{10}{3} + t\right]t\right)^8$ $t \text{ term: } \left(\frac{3}{16}\right)^8 \left[8 \times \frac{10}{3}\right] = 4.07 \times 10^{-5}$	M1 A1 M1M1 A1 [5]	Substitute and multiply Correctly obtain given answer Select t term; method for expansion formula Answer OR: attempt to find $G'(0)$ M1 $8\left(\frac{3}{16}\right)^8 \left(\frac{10}{3} + 2t\right) \left(1 + \left[\frac{10}{3} + t\right]t\right)^7$ A1 $= 4.07 \times 10^{-5}$ A1	

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2016	9795	02

4	(i)	Number of goals scored by home team is independent of number of goals scored by away team	B1 [1]	Not just <i>goals</i> independent. Extras, including conditions already implied by given Poisson distributions: B0
	(ii) (a)	$e^{-4.2n} \left(1 + 4.2n + \frac{(4.2n)^2}{2!} + \frac{(4.2n)^3}{3!} \right)$	M1 A1 A1 [3]	Po(4.2n) implied Correct ± 1 term Fully correct expression, aef. SR Po(4.2): Fully correct formula B1
	(b)	$e^{-2.4} e^{-1.8} (1 + 2.4 \times 1.8)$ $= \mathbf{0.0798}$	M1 A1 A1 [3]	Individual Poisson distributions multiplied Correct expression [= 0.0150 + 0.0647] Answer, a.r.t. 0.080 [0.07977]
5	(i)	n large p close to $\frac{1}{2}$	B1 B1 [2]	Or $np > 5$ $nq > 5$ [not $npq > 5$]
	(ii)	$\frac{24.5 - \mu}{\sigma} = \Phi^{-1}(0.8282) = 0.947$ $\frac{27.5 - \mu}{\sigma} = \Phi^{-1}(0.9697) = 1.759$ $\mu = \mathbf{21}, \sigma = \mathbf{3.69}$	M1 A1 B1 M1 A1 [5]	One standardised, = Φ^{-1} , allow σ^2 , cc, 1- errors LHS of both equations correct including signs and cc Both z -values correct to 3 sf, ± 1 in third dp Solve to find both μ and σ μ , a.r.t. 21.0; σ , in range [3.69, 3.70]
	(iii)	$q = npq/np = 21/3.694^2$ [= 0.65] $p = \mathbf{0.35}, n = \mathbf{60}$	M1 A1ft A1 [3]	Correct method of solution for n, p or q , allow \sqrt{npq} $npq = \sigma^2$ [not σ], ft on their npq [13.65] p , a.r.t. 0.350 and $n = 60$ [integer] <i>only</i> [not 60.0]
6	(i)	$\int_0^{\infty} 4xe^{-2x} e^{tx} dx = \int_0^{\infty} 4xe^{-(2-t)x} dx$ $= \left[\frac{4xe^{-(2-t)x}}{(t-2)} \right]_0^{\infty} + \int_0^{\infty} \frac{4e^{-(2-t)x}}{2-t} dx$ $\left[-\frac{4e^{-(2-t)x}}{(2-t)^2} \right]_0^{\infty} = \frac{4}{(2-t)^2}$	M1 A1 M1 A1 A1 [5]	Attempt $\int e^{tx} f(x) dx$, limits somewhere Combine into single e term Use parts, right way round Correct indefinite integral Correct final answer, cwo, allow $(t-2)^2$ but must use integral that visibly converges, or otherwise indicate the issue
	(ii)	$t < 2$	B1 [1]	
	(iii)	$\left[\frac{4}{(2-t)^2} \right]^3 = \frac{64}{(2-t)^6}$ $= (1 - \frac{1}{2}t)^{-6} = 1 + 3t + \frac{21}{4}t^2 + \dots$ $E(Y) = \mathbf{3}$ $E(Y^2)/2 = 21/4$ so $E(Y^2) = 10.5$ $\text{Var}(Y) = 10.5 - 3^2 = \mathbf{1.5}$	M1 A1 M1 A1 A1 [6]	$[M_X(t)]^3$ [Not cubed: M0A0 M1A0 M1A0] Series expansion <i>or</i> differentiate once $M'(t) = \frac{384}{(2-t)^7}, M''(t) = \frac{2688}{(2-t)^8}$ $E(Y) = 3$ correctly obtained or implied $2 \times \text{coeff of } t^2 \text{ or } M''(0) - [M'(0)]^2$ $\text{Var}(Y) = 1.5$ or exact equivalent, cwo

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2016	9795	02
7	<p>(i) $\int_0^k x \frac{3x^2}{k^3} dx = \frac{3}{4}k$ $E(\frac{4}{3}X) = k$, so $\frac{4}{3}X$ unbiased AG</p> <p>(ii) $P(X \leq x) = \int_0^k \frac{3x^2}{k^3} dx = \left(\frac{x^3}{k^3}\right)$ $P(M \leq m) = \left(\frac{x^3}{k^3}\right)^3 = \frac{x^9}{k^9}$ $f_M(x) = 9 \frac{x^8}{k^9}$ AG</p> <p>(iii) $\int_0^k x 9 \frac{x^8}{k^9} dx = \frac{9}{10}k$ Hence $E_2 = \frac{10}{9}M$</p>	<p>M1 A1 A1 [3]</p> <p>B1 M1 M1 A1 [4]</p> <p>M1 A1 A1ft [3]</p>	<p>Attempt $\int xf(x)$, correct limits $\frac{3}{4}k$, ae exact f Must state “unbiased”</p> <p>Needs convincing derivation $[F_X(x)]^3$ Differentiate Full derivation of AG. Ignore other ranges</p> <p>Attempt $\int xf_M(x)$, ignore limits Correct $E(M)$ If $E(M) = kc$, allow M/c</p>
8	<p>PE lost = $0.4g \times 3 \sin 20^\circ$ [4.104] Initial KE = $\frac{1}{2} \times 0.4 \times 0.5^2$ [0.05] Final KE = $\frac{1}{2} \times 0.4 \times 2.5^2$ [1.25] Difference = Work done by friction $2.9045 = 3F$ Therefore $F = \mathbf{0.968}$ N</p>	<p>M1 M1 M1 M1 A1 [5]</p>	<p>mgh attempted, with trig Both KEs attempted</p> <p>Work/Energy principle used, no extra/missing terms</p> <p>Use $\Delta E = F \times s$ Answer, a.r.t. 0.968 [SC: Energy not used: answer B2]</p>
9	<p>(i) $R(\uparrow_Q): T - 1.5g = 0$ [$T = 15$] $R(\uparrow_P): T \cos \theta - 1.2g = 0$ [$15 \cos \theta = 12$] $\Rightarrow \cos \theta = 0.8$ Distance below = $0.12/\tan \theta$ = 0.16m</p> <p>(ii) $T = 1.5g$ $R(\rightarrow_P): T \sin \theta = 1.2r\omega^2$ [= 9] $\Rightarrow 1.5g \sin \theta = 1.2 \times 0.12 \times \omega^2$ [$a = 7.5$] $\Rightarrow \omega = \sqrt{62.5} = \mathbf{7.91}$ rad s^{-1}</p>	<p>M1 M1 A1 A1 [4]</p> <p>B1 M1 A1 A1 [4]</p>	<p>Resolve vertically for 1.5 kg mass, can be implied e.g. by $T = 15$ Resolve vertically for 1.2 kg mass</p> <p>Value of $\cos \theta$ [$\theta = 36.9^\circ$] Correct value of h</p> <p>Value of T used [= 15] Resolve horiz for P and use $r\omega^2$ or v^2/r Correct equation, and $v = r\omega$ if v used</p> <p>Answer, in range [7.9, 7.91] or $\frac{5}{2}\sqrt{10}$</p>

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2016	9795	02
10 (i)	$N = 80$ $M(X): F \times 5 \sin \theta = 80 \times 2.5 \cos \theta$ $F \leq 0.4N$ $\tan \theta \geq 1.25$ $\theta_{\min} = 51.3^\circ$ or $51.4^\circ, 0.896$	B1 M1 M1 M1 A1	Normal force at ground (can be implied) Moments about any point, needs both cos and sin $A: R \times 5 \sin \theta = 80 \times 2.5 \cos \theta$ $[R = F = 0.4N = 32]$ $M: R \times 2.5 \sin \theta + F \times 2.5 \cos \theta = N \times 2.5 \cos \theta$ Use $F \leq \mu N$ or $F = \mu N$ Solve equations to obtain $\tan \theta$ Correct answer, in range $[51.3, 51.4]$ or a.r.t. 0.896
(ii)	$F \times 5 \sin \theta = (80 \times 2.5 + 750d) \cos \theta$ Use 60° and μ to obtain $d_{\max} = 3.57$	M1* depM1 A1	Moments equation with variable (d) [$F = 332$] $[332 \times 5 \times 0.5 = 100\sqrt{3} + 337\sqrt{3}d]$ Answer, a.r.t. 3.57 or 3.56
11 (i)	Driving force = $32000/v$ $800 \frac{dv}{dt} = \frac{32000}{v} - 20v$ $\frac{dv}{dt} = \frac{1600 - v^2}{40v}$ AG	M1 A1	Use P/v and differential equation including dv/dt Correctly obtain AG, need to use 800 convincingly
(ii)	$\int \frac{40v}{1600 - v^2} dv = \int dt$ $c - 20 \ln(1600 - v^2) = t$ $c = 20 \ln 1600$ [147.56] $t = 20 \ln \left(\frac{1600}{1600 - v^2} \right)$ $v = 40 \sqrt{1 - e^{-t/20}}$ Tends to 40	M1 A1 A1 M1 A1 B1	Separate variables and attempt to integrate Correct indefinite integral, aef Correct value of c Make v subject, using e , allow v^2 Correct expression for v , aef Conclusion, cwo but can get from implicit formula
12 (i)	 $\text{Mom}^m (\rightarrow): mu \cos 30 = mw + mx$ $\text{Rest}^n: x - w = 0.9\sqrt{3}u/2$ Solve: $w = 0.0433u$ $\text{Mom}^m (\uparrow): mu \sin 30 = vm$ so $v = 0.5u$ \Rightarrow direction is $\tan^{-1}(0.5/0.0433)$ $= 85.05^\circ$ to x -axis	M1 M1 A1 B1 A1	$[\frac{\sqrt{3}}{2}u = w + x]$ C of M equation, needn't have m , ignore signs, needs cos and sin Restitution equation, ignore signs of LHS Correctly obtain $w = \text{a.r.t. } 0.0433u$ [= $\sqrt{3}u/40$] Obtain, state, or use $v = u/2$ Direction, $[85.0^\circ, 85.1^\circ]$ to x -axis (5° or 4.9° to y)
(ii)	$u \cos 30 = w + x, ue \cos 30 = x - w$ $\Rightarrow 2w = u \cos 30(1 - e)$ but $e \leq 1$ so w cannot be negative	M1 M1 A1	One general equation Second equation, and use $e \leq 1$ Correctly deduce given conclusion

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2016	9795	02

<p>13 (i)</p>	<p>Solution 1: V, H</p> $y = 30t \sin 50^\circ - 5t^2; x = 30t \cos 50^\circ$ $[y = 26.2] \quad x = 48.2$ <p>Height of slope = $x \tan 10^\circ = 8.5$ Difference = 17.7</p> <p>Solution 2: \parallel, \perp</p> $y = 30t \sin 40^\circ - 5t^2 \cos 10^\circ$ $= 17.43$ <p>Height above field = $y \div \cos 10^\circ$ = 17.7</p>	<p>M1A1 A1</p> <p>M1 A1</p> <p>[5]</p>	<p>Both equations attempted; both correct Correct value of x SC: If M0, $y = 26.2$ gets B1</p> <p>Find height Correct answer [17.703]</p>
<p>(ii)</p>	<p>Solution 1: V, H</p> $30t \sin 50^\circ - 5t^2 = 30t \cos 50^\circ \tan 10^\circ$ $t = 3.916$ $PX = 30t \cos 50^\circ \div \cos 10^\circ$ $X = \mathbf{76.7}$ <p>Solution 2: \parallel, \perp</p> $Y = 30t \sin 40^\circ - 5t^2 \cos 10^\circ$ $Y = 0 \text{ at } t = 3.916$ $X = 30t \cos 40^\circ - 5t^2 \sin 10^\circ$ $PX = \mathbf{76.7}$ <p>Solution 3: traj</p> $Y = X \tan 50^\circ - \frac{10X^2 \sec^2 50^\circ}{2 \times 30^2}$ $= x \tan 10^\circ$ $PX = 180(\tan 50^\circ - \tan 10^\circ) \cos^2 50^\circ =$ $\mathbf{76.7}$	<p>M1 A1</p> <p>M1 A1</p> <p>[4]</p> <p>M1 A1</p> <p>A1 A1</p> <p>M1 A1</p>	<p>Put $y = x \tan 10^\circ$ and solve Correct value of t, can be implied Calculate $x \div \cos 10^\circ$ or $y \div \sin 10^\circ$ X, a.r.t. 76.7 [76.684]</p> <p>Y equation, allow sign/trig errors (3 terms) Correct value of t, can be implied X equation, allow sign/trig errors (3 terms) X, a.r.t. 76.7 [76.684]</p> <p>Use trajectory equation All correct</p> <p>Solve for X X, a.r.t. 76.7 [76.684]</p>

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2016	9795	02

14 (i)	$T - mg = 0 \Rightarrow \frac{3}{0.5} \times e = 0.3g$ $\Rightarrow e = 0.5$	M1	Use N2 for equilibrium	
	(ii)	GPE lost = EPE gained: $3(0.5 + x) = \frac{1}{2} \frac{3}{0.5} x^2 \quad [= 3x^2]$ Solve to get $x = \frac{1 + \sqrt{3}}{2} = 1.37$	A1 [2]	Equilibrium extension 0.5, a.r.t. 0.500
		M1*	Use cons of energy, need variable on both sides, e.g. $3(1 + h) = 3(h + \frac{1}{2})^2$ or $3h = 3(h - \frac{1}{2})^2$	
		depM1 A1 A1 [4]	Obtain and solve quadratic equation Correct equation, add/subtract 0.5 etc if necessary Solve to get a.r.t. 1.37 <i>only</i> , allow surds	
(iii)	$0.3\ddot{x} = 0.3g - \frac{3}{0.5}(x + e)$ $\ddot{x} = -20x$	M1 A1 B1 [3]	Use N2 including extension All correct, check signs Obtain correct value of ω^2 , no wrong working	
(iv)	$x = \frac{\sqrt{3}}{2} \cos \sqrt{20}t$ $x = 0.5$ at $\omega t = 0.9553$ or 5.3279 $t = 0.2136$ or 1.1913 Difference = 0.978	M1 A1ft M1 A1 A1 [5]	Use $x = a \cos \omega t$ or $a \sin \omega t$, allow $a = 1$ a correct ft (= their (ii) – their (i)), ω from (iii) Equate to $(\pm)e$ and use correct trig method e.g. $\frac{2}{\sqrt{20}} \cos^{-1} \left(-\frac{1}{\sqrt{3}} \right)$ or $\frac{1}{\sqrt{20}} \left(\frac{\pi}{\omega} + 2 \sin^{-1} \left(\frac{1}{\sqrt{3}} \right) \right)$ One correct value of t Correct final answer [0.9777]	