



**Cambridge International Examinations**  
Cambridge Pre-U Certificate

---

**FURTHER MATHEMATICS**

**9795/01**

Paper 1 Further Pure Mathematics

**May/June 2017**

MARK SCHEME

Maximum Mark: 120

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

© IGCSE is a registered trademark.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 3 Pre-U Certificate.

---

This document consists of **11** printed pages.

| Question | Answer   | Marks       | Part Marks   |
|----------|--|-------------|--|
| 1        | $(a + ib)^2 = (a^2 - b^2) + i.2ab$   | <b>B1</b>   |  |
|          | $(a^2 - b^2) = 21$ and $ab = -10$  | <b>M1</b>   | Comparing real and imaginary parts   |
|          | e.g. eliminating one variable and solving for the other  | <b>M1</b>   | Allow implied by e.g. $a = 5$ , $b = 2$ (or v.v.)  |
|          | $a = \pm 5$ , $b = \mp 2$  | <b>A1</b>   | Ignore any complex answers   |
| 2        | $\Sigma\alpha = -2$ and $\Sigma\alpha\beta = 3$  | <b>B1</b>   | Both ( $\alpha\beta\gamma = -7$ not required)  |
|          | $\alpha^2 + \beta^2 + \gamma^2 = (\Sigma\alpha)^2 - 2\Sigma\alpha\beta = -2$                                     | <b>M1A1</b> | <b>FT</b>  |
|          | 1 real and 2 complex (conjugate) roots   | <b>B1</b>   | Accept any comment that “not all roots are real  |
|          | <b>Alternative</b><br>Form an equation with roots $\alpha^2, \beta^2, \gamma^2$ ;<br>$y^3 + 2y^2 - 19y - 49 = 0$ | <b>M1A1</b> |  |
|          | $\Sigma\alpha^2 = -\frac{b}{a} = -2$   | <b>B1</b>   | <b>FT</b>  |
|          | 1 real and 2 complex (conjugate) roots   | <b>B1</b>   | Accept any comment that “not all roots are real  |
| 3(i)     |  | <b>B3</b>   | <b>B1</b> Starts at (1, 0)<br><b>B1</b> Decreasing spiral<br><b>B1</b> All (essentially) correct |
| 3(ii)    | $\text{Area} = \frac{1}{2} \int_0^{2\pi} \frac{1}{(1+\theta)^2} d\theta$   | <b>M1</b>   | Attempt to integrate $k(1+\theta)^{-2}$  |
|          | $= \frac{1}{2} \left[ \frac{-1}{1+\theta} \right]_0^{2\pi}$  | <b>A1</b>   | Correct integration  |
|          | $= \frac{1}{2} \left( 1 - \frac{1}{1+2\pi} \right)$ or $\frac{\pi}{1+2\pi}$                                      | <b>A1</b>   | Correct answer   |

| Question | Answer   | Marks     | Part Marks   |
|----------|--|-----------|--|
| 4        | $\dot{x} = t - \frac{1}{t}$ and $\dot{y} = 2$  | <b>B1</b> | at least $\dot{x}$ correct   |
|          | $(\dot{x})^2 + (\dot{y})^2 = t^2 - 2 + \frac{1}{t^2} + 4$  | <b>M1</b> | attempted  |
|          | $= \left(t + \frac{1}{t}\right)^2$   | <b>A1</b> | Here or in the integral for $S$ (2 <sup>nd</sup> fraction of line below)                       |
|          | $S = 2\pi \int_1^4 2t \cdot \left(t + \frac{1}{t}\right) dt$   | <b>M1</b> | Use of formula (Ignore limits until final answer)  |
|          | $= 4\pi \int_1^4 (t^2 + 1) dt$   | <b>A1</b> | In a form ready to integrate   |
|          | $= 4\pi \left[ \frac{t^3}{3} + t \right]_1^4$  | <b>B1</b> | Correct integration ( <b>FT</b> provided it is polynomial)                                     |
|          | $= 96\pi$  | <b>A1</b> |  |
| 5(i)     | $y = \tanh^{-1} x \Leftrightarrow \tanh y = x = \frac{e^{2y} - 1}{e^{2y} + 1}$   | <b>M1</b> |  |
|          | $xe^{2y} + x = e^{2y} - 1 \Leftrightarrow 1 + x = e^{2y}(1 - x)$   | <b>M1</b> | Identifying $e^{2y}$   |
|          | $y = \tanh^{-1} x = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$  | <b>A1</b> | Legitimately obtained by taking logs<br><br>Allow verification by substitution of given result |
| 5(ii)    | <b>Method I</b><br>$t + \frac{1}{t} = 4 \Rightarrow t^2 - 4t + 1 = 0$  | <b>M1</b> | Creating a quadratic in $\tanh x$  |
|          | $\Rightarrow t = 2 \pm \sqrt{3}$   | <b>M1</b> | Solving  |
|          | Using $\frac{1}{2} \ln \left( \frac{1+t}{1-t} \right)$ with $t = 2 - \sqrt{3}$ and/or $2 + \sqrt{3}$   | <b>M1</b> | (NB since $ \tanh x  < 1$ , it must be $t = 2 - \sqrt{3}$ )                                    |
|          | $x = \frac{1}{2} \ln \left( \frac{3 - \sqrt{3}}{-1 + \sqrt{3}} \times \frac{1 + \sqrt{3}}{1 + \sqrt{3}} \right) = \frac{1}{2} \ln(\sqrt{3})$ | <b>M1</b> | By rationalising denominator or direct observation (possibly from calculator use)              |
|          | $= \frac{1}{4} \ln(3)$   | <b>A1</b> | Must be in this form   |

| Question | Answer  | Marks       | Part Marks   |
|----------|---|-------------|--|
| 5(ii)    | <b>Method II</b><br>$\frac{\text{sh}}{\text{ch}} + \frac{\text{ch}}{\text{sh}} = 4$   | <b>M1</b>   |  |
|          | $\Rightarrow \text{ch}^2 + \text{sh}^2 = 4\text{sh.ch} \Rightarrow \cosh(2x) = 2 \sinh(2x)$   | <b>M1</b>   | Conversion to double-“angles”  |
|          | $\Rightarrow \tanh(2x) = \frac{1}{2}$   | <b>A1</b>   |  |
|          | $\Rightarrow 2x = \frac{1}{2} \ln\left(\frac{\frac{3}{2}}{\frac{1}{2}}\right)$  | <b>M1</b>   | Use of $\tanh^{-1} x$ formula from (i)   |
|          | $\Rightarrow x = \frac{1}{4} \ln(3)$  | <b>A1</b>   | Must be in this form   |
|          | <b>Method III</b><br>$\frac{e^{2x}-1}{e^{2x}+1} + \frac{e^{2x}+1}{e^{2x}-1} = 4$  | <b>M1</b>   |  |
|          | $\Rightarrow (e^{2x}-1)^2 + (e^{2x}+1)^2 = 4(e^{2x}-1)(e^{2x}+1)$   | <b>M1</b>   |  |
|          | $\Rightarrow e^{4x} - 2e^{2x} + 1 + e^{4x} + 2e^{2x} + 1 = 4(e^{4x} - 1)$   | <b>A2</b>   | <b>A1</b> LHS<br><b>A1</b> RHS   |
|          | $\Rightarrow 6 = 2e^{4x} \Rightarrow x = \frac{1}{4} \ln(3)$  | <b>A1</b>   | Must be in this form   |
| 6(i)     | HA $y = 1$ VA $x = -1$  | <b>B2</b>   | <b>B1</b> for each   |
| 6(ii)    | $y = \frac{x^2+1}{(x+1)^2}$ or $y = 1 - \frac{2x}{(x+1)^2}$<br>$\Rightarrow \frac{dy}{dx} = \frac{(x+1)^2(2x) - (x^2+1).2(x+1)}{(x+1)^4}$ or<br>$-\frac{(x+1)^2.2 - 2x.2(x+1)}{(x+1)^4} = \frac{2(x-1)}{(x+1)^3}$ | <b>M1A1</b> | Attempted; correct unsimplified  |
|          | $\Rightarrow \frac{dy}{dx} = 0$ when $x = 1, y = \frac{1}{2}$   | <b>A2</b>   | <b>A1</b> for each   |
| 6(iii)   |   | <b>3</b>    | <b>G1</b> for graph in 2 bits, separated by a (FT) vertical asymptote and all positive<br><br><b>G1</b> for y-intercept at (0, 1) and MIN. in (approx. FT) correct place<br><br><b>G1</b> for correct asymptotic behaviour |

| Question | Answer  | Marks       | Part Marks  |
|----------|---|-------------|---|
| 7(i)     | $y = kx \sin 2x \Rightarrow \frac{dy}{dx} = 2kx \cos 2x + k \sin 2x$  | <b>M1</b>   | attempt using the <i>Product Rule</i>   |
|          | and $\frac{d^2y}{dx^2} = kx \cdot -4 \sin 2x + 2k \cos 2x + 2k \cos 2x$   | <b>M1</b>   | attempt using the <i>Product Rule</i>   |
|          | $= -4y + 4k \cos 2x$  | <b>M1</b>   | for substn. into given d.e. or comparison   |
|          | $\Rightarrow k = 2$   | <b>A1</b>   |   |
| 7(ii)    | Comp. Fn. from $m^2 + 4 = 0$  | <b>M1</b>   |   |
|          | $\Rightarrow y_c = A \cos 2x + B \sin 2x$   | <b>A1</b>   | Or $R \cos(2x - \alpha)$ etc.   |
|          | Gen. Soln. is thus $y = A \cos 2x + (B + 2x) \sin 2x$   | <b>B1</b>   | <b>FT</b>   |
|          | Then $\frac{dy}{dx} = -2A \sin 2x + 2(B + 2x) \cos 2x + 2 \sin 2x$<br>OR $= 2(B + 2x) \cos 2x$ if found after $A$ (correctly) evaluated | <b>B1</b>   |   |
|          | Subst <sup>g</sup> . in given initial conditions  | <b>M1</b>   |   |
|          | $A = 1$ from $x = 0, y = 1$   | <b>A1</b>   | <b>FT</b> from an incorrect $x \sin 2x$ term in $y$   |
|          | $B = \frac{1}{2}$ from $x = 0, \frac{dy}{dx} = 1$<br>i.e. soln. is $y = \cos 2x + (2x + \frac{1}{2}) \sin 2x$                           | <b>A1</b>   | <b>FT</b> from an incorrect $x \cos 2x$ term in $y'$<br><br>Withhold final A mark if in $e^{\wedge}$ complex form                     |
| 8(i)(a)  | $\cos \theta = \frac{12 + 2 + 6}{3 \times 7} = \frac{20}{21}$   | <b>M1A2</b> | <b>A1</b> scalar product; <b>A1</b> both moduli<br><br>Give B1s for correct scalar product; both moduli if $\sin \theta = \dots$ used |
| 8(i)(b)  | Subst <sup>g</sup> . $(2\lambda, -\lambda, 2\lambda)$ into $6x - 2y + 3z = 35$  | <b>M1</b>   |   |
|          | $\Rightarrow \lambda = \frac{7}{4} \Rightarrow \mathbf{p} = \frac{7}{4} \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$                     | <b>A1A1</b> | Second <b>A1</b> is <b>FT</b>   |
| 8(i)(c)  | SD $O$ to $\Pi_1 = OP \cos \theta = \frac{7}{4} \times 3 \times \frac{20}{21} = 5$  | <b>M1A1</b> | <b>A1FT</b>   |

| Question | Answer  | Marks       | Part Marks  |
|----------|---|-------------|---|
| 8(i)(c)  | <b>Alternative I</b><br>$(6\lambda, -2\lambda, 3\lambda)$ in plane $\Rightarrow 36\lambda + 4\lambda + 9\lambda = 35$   | <b>M1</b>   | $\Rightarrow \lambda = \frac{5}{7}$                                   |
|          | $\Rightarrow SD = \lambda \sqrt{6^2 + 2^2 + 3^2} = 5$ cao   | <b>A1</b>   |   |
|          | <b>Alternative II</b><br>Quote formula: $SD = \frac{\left  \frac{d}{\ \mathbf{n}\ } \right }{\text{cao}} = \frac{35}{\sqrt{6^2 + 2^2 + 3^2}} = 5$                             | <b>M1A1</b> |   |
| 8(ii)    | Similar working gives $\lambda_1 = -\frac{21}{40}$  | <b>B1</b>   |   |
|          | Planes parallel, <i>and on opposite sides of O</i> ,<br>so total distance is $3\left(\frac{7}{4} + \frac{21}{40}\right) \cos \theta = \frac{13}{2}$                           | <b>M1A1</b> |   |
|          | <b>Alternative I</b><br>$\Pi_2$ has equation $\mathbf{r} \cdot \begin{pmatrix} 6 \\ -2 \\ 3 \end{pmatrix} = -\frac{21}{2}$  | <b>B1</b>   |   |
|          | $\Rightarrow SD$ to $\Pi_2$ is $-\frac{3}{2}$   | <b>B1</b>   |   |
|          | Planes parallel, <i>and on opposite sides of O</i> ,<br>so distance between them is $5 - -\frac{3}{2} = \frac{13}{2}$   | <b>B1</b>   | <b>FT</b>   |
|          | <b>Alternative II</b><br>Quote Sh. Dist. formula for $P\left(\frac{7}{4}, -\frac{7}{2}, \frac{7}{2}\right)$ to $\Pi_2$  | <b>M1</b>   | or using distance from any point in $\Pi_1$ or $\Pi_2$ to other plane |
|          | $SD = \frac{\left  12\left(\frac{7}{4}\right) - 4\left(-\frac{7}{2}\right) + 6\left(\frac{7}{4}\right) + 21 \right }{\sqrt{12^2 + 4^2 + 6^2}} = \frac{91}{14} = \frac{13}{2}$ | <b>A1A1</b> |   |
| 9(i)     | Full elimination of $x$ : $I = \int \frac{1}{\cosh^2 \theta \cdot \sinh \theta} \cdot \sinh \theta d\theta$   | <b>M1</b>   |   |
|          | $\Rightarrow I = \int \text{sech}^2 \theta d\theta$   | <b>A1</b>   |   |
|          | $= \tanh \theta (+ C)$  | <b>A1</b>   |   |
|          | $= \frac{\sqrt{x^2 - 1}}{x} (+ C)$ from $\frac{\sinh \theta}{\cosh \theta}$   | <b>A1</b>   | <b>(AG)</b>   |

| Question | Answer  | Marks       | Part Marks   |
|----------|---|-------------|--|
| 9(ii)    | $\sec y = x \Rightarrow \sec y \tan y \frac{dy}{dx} = 1$  | <b>M1A1</b> |  |
|          | Use of $\tan y = \sqrt{\sec^2 y - 1}$   | <b>M1</b>   |  |
|          | to get $\frac{dy}{dx} = \frac{1}{x\sqrt{x^2 - 1}}$  | <b>A1</b>   | <b>AG</b><br>Ignore lack of reason for taking the +ve sq.rt. (e.g. from +ve gradient of $\sec^{-1}$ curve) |
| 9(iii)   | $\int \sec^{-1} x \cdot \frac{1}{x^2} dx$<br>$= \sec^{-1} x \cdot \frac{-1}{x} - \int \frac{-1}{x} \cdot \frac{1}{x\sqrt{x^2 - 1}} dx$<br>$= \frac{-\sec^{-1} x}{x} + \int \frac{1}{x^2 \sqrt{x^2 - 1}} dx$ | <b>M1A2</b> | By parts   |
|          | $= \frac{-\sec^{-1} x}{x} + \frac{\sqrt{x^2 - 1}}{x} (+C)$  | <b>A1</b>   | using (i)  |
|          | <b>Alternative</b><br>Use $u = \sec^{-1} x \Rightarrow \frac{du}{dx} = \frac{1}{x\sqrt{x^2 - 1}}$<br>$\Rightarrow \sec u \tan u du = dx$  | <b>M1</b>   |  |
|          | $\Rightarrow \int \sec^{-1} x \cdot \frac{1}{x^2} dx = \int u \sin u du$  | <b>A1</b>   |  |
|          | 2-stage integration by parts:<br>$\int u \sin u du = -u \cos u + \int \cos u du$<br>$= -u \cos u + \sin u (+C)$   | <b>M1</b>   |  |
|          | Correctly turning this back into<br>$= \frac{-\sec^{-1} x}{x} + \frac{\sqrt{x^2 - 1}}{x} (+C)$  | <b>A1</b>   |  |
| 10(i)    | $\frac{1}{(k-1)k(k+1)} \equiv \frac{A}{k-1} + \frac{B}{k} + \frac{C}{k+1}$  | <b>M1</b>   | Correct form   |
|          | Equating terms / substn. / cover-up   | <b>M1</b>   | Method for determining constants   |
|          | $\equiv \frac{\frac{1}{2}}{k-1} - \frac{1}{k} + \frac{\frac{1}{2}}{k+1}$  | <b>A1</b>   |  |

| Question | Answer  | Marks     | Part Marks                       |
|----------|---|-----------|----------------------------------|
| 10(ii)   | $\sum_{k=3}^n \frac{1}{(k-1)k(k+1)} \equiv \frac{1}{2} \sum_{k=3}^n \frac{1}{k-1} + \frac{1}{2} \sum_{k=3}^n \frac{1}{k+1} - \sum_{k=3}^n \frac{1}{k}$  | <b>M1</b> | Splitting up                     |
|          | $\equiv \frac{1}{2} \left\{ \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n-1} \right\} + \frac{1}{2} \left\{ \frac{1}{4} + \dots + \frac{1}{n-1} + \frac{1}{n} + \frac{1}{n+1} \right\} - \left\{ \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n-1} + \frac{1}{n} \right\}$ | <b>M1</b> | Attempt at cancelling of terms   |
|          | $\equiv \frac{1}{2} \left\{ \frac{1}{2} + \frac{1}{3} \right\} + \frac{1}{2} \left\{ \frac{1}{n} + \frac{1}{n+1} \right\} - \left\{ \frac{1}{3} + \frac{1}{n} \right\}$   | <b>A1</b> | Correct ones clearly identified  |
|          | $\equiv \frac{1}{12} - \frac{1}{2} \left\{ \frac{1}{n} - \frac{1}{n+1} \right\} \equiv \frac{1}{12} - \frac{1}{2n(n+1)}$  | <b>A1</b> | Legitimately shown ( <b>AG</b> ) |
|          | Limit ( $S_n$ ) as $n \rightarrow \infty$ is $S = \frac{1}{12}$   | <b>B1</b> | <b>FT</b>                        |
|          | <b>Alternative</b><br>$\sum_{k=3}^n \frac{1}{(k-1)k(k+1)} \equiv \frac{1}{2} \sum_{k=3}^n \frac{1}{k(k-1)} - \frac{1}{2} \sum_{k=3}^n \frac{1}{k(k+1)}$   | <b>M1</b> |                                  |
|          | $= \frac{1}{2} \left( \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \dots + \frac{1}{n(n-1)} \right) - \frac{1}{2} \left( \frac{1}{12} + \frac{1}{20} + \dots + \frac{1}{n(n-1)} + \frac{1}{n(n+1)} \right)$  | <b>M1</b> | Clear listing of terms           |
|          | All correct and ready to cancel   | <b>A1</b> |                                  |
|          | $= \frac{1}{12} - \frac{1}{2n(n+1)}$  | <b>A1</b> | Legitimately shown ( <b>AG</b> ) |
|          | Limit ( $S_n$ ) as $n \rightarrow \infty$ is $S = \frac{1}{12}$   | <b>B1</b> | <b>FT</b>                        |
| 10(iii)  | $k^3 > k^3 - k = k(k-1)(k+1)$<br>$\Rightarrow \frac{1}{k^3} < \frac{1}{(k-1)k(k+1)}$  | <b>B1</b> |                                  |
| 10(iv)   | $\sum_{k=1}^{\infty} \frac{1}{k^3} > 1 + \frac{1}{8} = \frac{9}{8} = \frac{27}{24}$   | <b>B1</b> | Given result justified           |
|          | $\sum_{k=1}^{\infty} \frac{1}{k^3} = 1 + \frac{1}{8} + \sum_{k=3}^{\infty} \frac{1}{k^3} < 1 + \frac{1}{8} + \sum_{k=3}^n \frac{1}{(k-1)k(k+1)}$  | <b>M1</b> |                                  |
|          | $= 1 + \frac{1}{8} + \frac{1}{12} = \frac{29}{24}$  | <b>A1</b> | Given result justified           |
| 11(i)(a) | $\mathbf{AB} = \begin{pmatrix} ae + bg & af + bh \\ ce + dg & cf + dh \end{pmatrix}$  | <b>B1</b> |                                  |
|          | $\det \mathbf{A} = ad - bc \text{ and } \det \mathbf{B} = eh - fg$  | <b>B1</b> |                                  |



| Question   | Answer   | Marks     | Part Marks  |
|------------|--|-----------|---|
| 11(i)(b)   | $\det(\mathbf{AB}) = (ae + bg)(cf + dh) - (af + bh)(ce + dg)$<br>and some attempt to multiply out  | <b>M1</b> |   |
|            | $= acef + adeh + bcfg + bdgh$<br>$\quad - acef - bceh - adfg - bdgh$<br>$= adeh - bceh - adfg + bcfg$<br>$= (ad - bc)(eh - fg)$  | <b>A1</b> | Legitimately shown  |
| 11(ii)     | <i>CLOSURE</i> : $\mathbf{A}, \mathbf{B} \in S \Rightarrow \det \mathbf{A} = \det \mathbf{B} = 1$  | <b>M1</b> | Attempted   |
|            | and above result $\Rightarrow \det \mathbf{AB} = 1 \Rightarrow \mathbf{AB} \in S$<br><i>(ASSOCIATIVITY: given)</i>   | <b>A1</b> | Convincing  |
|            | <i>IDENTITY</i> : $\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \in S$<br>since $\det \mathbf{I} = 1.1 - 0.0 = 1$   | <b>B1</b> | Must show why $\mathbf{I} \in S$ and not just say that $\mathbf{I}$ is the identity |
|            | <i>INVERSES</i> : $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in S \Rightarrow \mathbf{A}^{-1}$<br>$= \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \in S$                                  | <b>B1</b> | for stating $\mathbf{A}^{-1}$ (or explaining that it exists)                        |
|            | Since $da - (-b)(-c) = ad - bc = 1$<br>Hence $(S, \times_M)$ is a group, $G$ .   | <b>B1</b> | for justifying its membership of $S$  |
| 11(iii)(a) | $\det \mathbf{K} = 1.0 - i.i = -i^2 = 1$ (so $\mathbf{K} \in S$ )  | <b>B1</b> |   |
| 11(iii)(b) | Attempt at powers of $\mathbf{K}$ ; $\mathbf{K}^2$ & $\mathbf{K}^3$  | <b>M1</b> |   |
|            | $\mathbf{K}^2 = \begin{pmatrix} 0 & i \\ i & -1 \end{pmatrix}$ and $\mathbf{K}^3 = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$   | <b>A1</b> |   |
|            | NB $\mathbf{K}^4 = \begin{pmatrix} -1 & -i \\ -i & 0 \end{pmatrix}$ and $\mathbf{K}^5 = \begin{pmatrix} 0 & -i \\ -i & 1 \end{pmatrix}$<br>$\Rightarrow \mathbf{K}^6 = \mathbf{I}$ and $H$ has order $n = 6$ | <b>A1</b> |   |
| 11(iii)(c) | e.g. The set of rotations about $O$ through multiples of $60^\circ$<br><br>OR $(\mathbf{K}^*) =$ group generated by $\begin{pmatrix} 1 & -i \\ -i & 0 \end{pmatrix}$   | <b>B1</b> | <b>FT</b> for any $n$   |
|            | Justifying the two are isomorphic  | <b>B1</b> | e.g. stating both are cyclic, etc.  |

| Question  | Answer   | Marks     | Part Marks   |
|-----------|--|-----------|--|
| 12(i)     | <b>Method I</b><br>$F_{n+2}(\theta) - \frac{1}{4} \sin^2(2\theta) F_{n+1}(\theta)$ $\equiv (c^2 + s^2)(c^{2n+4} + s^{2n+4})$ $- \frac{1}{4}(2sc)^2(c^{2n+2} + s^{2n+2})$         | <b>M2</b> | <b>M1</b> all $F_n$ terms<br><b>M1</b> $\sin 2\theta$ form |
|           | $\equiv c^{2n+6} + c^2 s^{2n+4} + s^2 c^{2n+4} + s^{2n+6}$ $- c^2 s^2(c^{2n+2} + s^{2n+2})$  | <b>A1</b> |  |
|           | $\equiv c^{2n+6} + s^{2n+6} \equiv F_{n+3}(\theta)$  | <b>A1</b> | <b>AG</b>  |
|           | <b>Method II</b><br>$\equiv c^{2n+4} + s^{2n+4} - s^2 c^2(c^{2n+2} + s^{2n+2})$  | <b>M1</b> | Use of $\sin 2\theta$ form                                 |
|           | $\equiv c^{2n+4} + s^{2n+4} - s^2 c^{2n+4} - c^2 s^{2n+4}$   | <b>A1</b> |  |
|           | $\equiv (1-s^2)c^{2n+4} + (1-c^2)s^{2n+4}$   | <b>M1</b> |  |
|           | $\equiv c^{2n+6} + s^{2n+6} \equiv F_{n+3}(\theta)$  | <b>A1</b> | <b>AG</b>  |
| 12(ii)(a) | Use of $z = c + is$ and $z^{-1} = c - is$  | <b>M1</b> |  |
|           | $z + z^{-1} = 2c$ and $z - z^{-1} = 2is$   | <b>A2</b> | <b>A1</b> for each   |
| 12(ii)(b) | <b>Method I</b><br>$(2c)^6 = (z + z^{-1})^6 = z^6 + 6z^4 + 15z^2 + 20$ $+ 15z^{-2} + 6z^{-4} + z^{-6}$   | <b>M1</b> |  |
|           | $= 2\cos 6\theta + 12\cos 4\theta + 30\cos 2\theta + 20$   | <b>A1</b> |  |
|           | $-(2s)^6 = (z - z^{-1})^6 = z^6 - 6z^4 + 15z^2 - 20$ $+ 15z^{-2} - 6z^{-4} + z^{-6}$ $= 2\cos 6\theta - 12\cos 4\theta + 30\cos 2\theta - 20$                                    | <b>B1</b> | <b>FT</b> (Must have – sign)                               |
|           | Subtracting:<br>$64(c^6 + s^6) = 12(z^4 + z^{-4}) + 40$ $= 12 \cdot 2\cos 4\theta + 40$  | <b>M1</b> |  |
|           | Dividing by 8: $8(c^6 + s^6) = 3\cos 4\theta + 5$  | <b>A1</b> | <b>AG</b>  |
|           | Use of $\cos 4\theta = 2\cos^2 2\theta - 1$ and $1 = \cos^2 2\theta + \sin^2 2\theta$  | <b>M1</b> |  |
|           | $\Rightarrow c^6 + s^6 = \frac{3}{8}(2\cos^2 2\theta) + \left(-\frac{3}{8} + \frac{5}{8}\right)(\cos^2 2\theta + \sin^2 2\theta)$ $= \cos^2 2\theta + \frac{1}{4}\sin^2 2\theta$ | <b>A1</b> | <b>AG</b>  |

| Question  | Answer  | Marks       | Part Marks  |
|-----------|---|-------------|---|
| 12(ii)(b) | <b>Method II</b><br>$\cos 4\theta = \operatorname{Re}(c + is)^4$  | <b>M1</b>   |   |
|           | $= c^4 - 6c^2s^2 + s^4 = c^4 - 6c^2(1-c^2) + (1-c^2)^2$<br>$= 8c^4 - 8c^2 + 1$  | <b>A1</b>   |   |
|           | $c^6 + s^6 = c^6 + (1-c^2)^3 = c^6 + 1 - 3c^2 + 3c^4 - c^6$   | <b>M1</b>   |   |
|           | $= 3c^4 - 3c^2 + 1$   | <b>A1</b>   |   |
|           | so that $8(c^6 + s^6) = 3\cos 4\theta + 5$  | <b>A1</b>   | <b>AG</b>   |
|           | Use of $\cos 4\theta = \cos^2 2\theta - \sin^2 2\theta$<br>and $1 = \cos^2 2\theta + \sin^2 2\theta$  | <b>M1</b>   |   |
|           | $\Rightarrow 8(c^6 + s^6) = 3\cos 4\theta + 5$<br>$= 3(\cos^2 2\theta - \sin^2 2\theta) + 5(\cos^2 2\theta + \sin^2 2\theta)$<br>$\Rightarrow c^6 + s^6 = \cos^2 2\theta + \frac{1}{4}\sin^2 2\theta$ | <b>A1</b>   | <b>AG</b>   |
| 12(iii)   | Case for $n = 1$ established in (ii) (b):   | <b>B1</b>   | noted explicitly (possibly at end)                          |
|           | Assume $c^{2k+4} + s^{2k+4} \leq \cos^2 2\theta + \frac{1}{2^{k+1}}\sin^2 2\theta$  | <b>B1</b>   | i.e. the case for $n = k$                                   |
|           | A clear statement of the result must be given,<br>possibly within what follows<br>Then $c^{2k+6} + s^{2k+6} =$<br>$c^{2k+4} + s^{2k+4} - \frac{1}{4}\sin^2 2\theta(c^{2k+2} + s^{2k+2})$              | <b>M1</b>   | attempt at $n = k + 1$ case using (i)'s identity            |
|           | $\leq \cos^2 2\theta + \frac{1}{2^{k+1}}\sin^2 2\theta - \frac{1}{4}\sin^2 2\theta(c^{2k+2} + s^{2k+2})$  | <b>M1</b>   | use of the induction hypothesis (i.e. the $n = k$ case)     |
|           | $= \cos^2 2\theta + \frac{1}{2^{k+2}}\sin^2 2\theta - \frac{1}{4}\sin^2 2\theta\left(c^{2k+2} + s^{2k+2} - \frac{1}{2^k}\right)$  | <b>M1A1</b> | splitting up the $\sin^2 2\theta$ term into two equal parts |
|           | $\leq \cos^2 2\theta + \frac{1}{2^{k+2}}\sin^2 2\theta$<br>Proof follows by induction since $\sin^2 2\theta \geq 0$ and<br>given result that $c^{2k+2} + s^{2k+2} \geq \frac{1}{2^k}$                 | <b>A1</b>   |   |