

Mark Scheme (Results)

Summer 2017

Pearson Edexcel International GCSE In Mathematics B (4MB0) Paper 01R



Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.edexcel.com, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2017
Publications Code 4MB0_01R_1706_MS
All the material in this publication is copyright
© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

• Types of mark

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- o cao correct answer only
- ft follow through
- o isw ignore subsequent working
- SC special case
- o oe or equivalent (and appropriate)
- o dep dependent
- o indep independent
- o awrt answer which rounds to
- eeoo each error or omission

No working

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

• Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

1 (a) (b)	57900	1 1	B1 B1	
(b)	0.05	1	B1	
				Total 2 marks
2 3200000 or 32×10 ⁵		2	M1	
	3.2×10 ⁶		A1	
				Total 2 marks
	•			
3 $(3x =)7 - 2$		2	M1	
	$\frac{5}{3}$ (oe)		A1 1	$1\frac{2}{3}$, 1. $\dot{6}$, 1. $\overline{6}$, awrt 1.67
	$\frac{3}{3}$			NB do not allow 1.66
				Total 2 marks
$\frac{14}{30} \times 360 \text{ (oe)}$		2	M1 f	full method required
30 ~ 300 (00)				
	168		A1	
				Total 2 marks
			T	
$\pm \frac{96-90}{96}$ or $\pm \frac{6}{96}$ or $\frac{90}{96} \times 100$		3	M1	
			M1 DEP	
	$6\frac{1}{4}$ (oe)		A1 6	$6.25, \frac{25}{4}$
NB -6.25 score M1M1A0				Total 3 marks

6	(a) (i)		$x \ge -5$	2	B1	
	(ii)		x < 6		B1	
	(b)	Single Line •	0 6	1	B1	FT their inequalities
		-5				allow -5 [
						Total 3 marks
			,		_	
7		$\Delta = 5 \times -2 - 4 \times -3 (= 2)$		3	M1	
		$1(-2 \ 3)$			M1	(Indep) allow $\Delta=1$
		$ \frac{1}{\Delta} \begin{pmatrix} -2 & 3 \\ -4 & 5 \end{pmatrix} $				
			$\begin{pmatrix} -1 & 1\frac{1}{2} \\ -2 & 2\frac{1}{2} \end{pmatrix} (oe)$		A1	Accept $\frac{1}{2} \begin{pmatrix} -2 & 3 \\ -4 & 5 \end{pmatrix}$
						Total 3 marks
8		$2\sqrt{3}-2\sqrt{2}+\sqrt{6}\times\sqrt{3}-\sqrt{6}\times\sqrt{2}$ or		3		Correctly expand brackets.
		$2\sqrt{3} - 2\sqrt{2} + \sqrt{18} - \sqrt{12}$				Allow one sign error.
		$\sqrt{18} = 3\sqrt{2}$ and $\sqrt{12} = 2\sqrt{3}$ or			M1	May be embedded within working
		$\sqrt{6} \times \sqrt{3} = 3\sqrt{2}$ and $\sqrt{6} \times \sqrt{2} = 2\sqrt{3}$			(DEP)	
		$2\sqrt{3} - 2\sqrt{2} + 3\sqrt{2} - 2\sqrt{3}$	$\sqrt{2}$		A1	dep on M2
			V -			cso
		NB Answers derived from decimal expansions score	M0M0A0			Total 3 marks

9	tangent drawn at (2, 2)		3	M1 Must only touch curve at (2,2)
	y increment x increment			M1(DEP) Attempt to find gradient for their tangent.
	* Increment	3		A1 dep on tangent drawn ft their tangent providing $2.5 \le \text{gradient} \le 4$
				Total 3 marks
10	Shaded area = $(2r)^2 - \pi r^2$, 0.858 r^2 or better. OR $\frac{\pi(r^2)}{4(r^2)} \times 100$		3	M1
	OR $\frac{\pi(r^2)}{4(r^2)} \times 100$ $\frac{4r^2 - \pi r^2}{4r^2} \times 100\% \text{ OR } 100 - \frac{\pi(r^2)}{4(r^2)} \times 100$			M1 DEP
		21.5		A1 awrt 21.4 or 21.5
				Total 3 marks
11 (a)	E	Q, Z, N correctly labelled	1	B1
(b)	$Q \qquad Z \qquad $	$\{4, \sqrt{2}, -3, \frac{2}{3}\}$ four elements in correct position	2	B2 or B1 for two in correct position Do not allow repeated elements in different areas
				Total 3 marks

12	(10 9	1	M1	Correct dimension and at least one
	13 9		A1	correct entry.
	-4 -10)	A1	Any three correct entries. All correct.
	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	/	AI	
	NB for a transcription error allow at most one value changed or two v	alues transposed.		Total 3 marks
13 (a)		1	B1	Correct region shaded.
(b)		1	B1	Correct region shaded.
(c)	$(F \cup G) \cap H$	(oe) 1	B1	Accept $(F \cap H') \cup (G \cap H')$ or $(F \cap G' \cap H') \cup (F \cap G \cap H') \cup (F' \cap G \cap H')$
			1	Do not accept $F \cup G \cap H'$
				Total 3 marks

14	$\frac{6}{10} \times \frac{4}{10} \text{ or } \frac{4}{10} \times \frac{6}{9} \text{ or } \frac{6}{10} \times \frac{6}{10} + \frac{4}{10} \times \frac{3}{9}$		3	M1
	$\frac{6}{10} \times \frac{4}{10} + \frac{4}{10} \times \frac{6}{9} \text{ or } 1 - \left(\frac{6}{10} \times \frac{6}{10} + \frac{4}{10} \times \frac{3}{9}\right)$			M1 DEP
		$\frac{38}{75}$ (oe)		A1 AWRT 0.507
				Total 3 marks
	,		<u>r</u>	
15	SF = $\sqrt[3]{\frac{1350}{400}} \left(= \frac{3}{2} \right)$ or $\sqrt[3]{\frac{400}{1350}} \left(= \frac{2}{3} \right)$ or $\sqrt[3]{1350}$: $\sqrt[3]{400}$ or 3:2 (oe)		3	M1 Obtain linear scale factor. This mark not awarded unless cube root taken at some stage.
	$360 \times \left(\sqrt[3]{\frac{1350}{400}}\right)^2$ or 360×2.25 or			M1(DEP)
	$360 \div \left(\sqrt[3]{\frac{400}{1350}}\right)^2 \text{ or } 360 \div \frac{4}{9}$			
		810		A1
				Total 3 marks
16	(2 1)(5) (2 1)(1) [1 2	M1 fordering growth
16	$\frac{(2x-1)(x+5)}{3(2x-1)(2x+1)} \text{ or } \frac{(2x-1)(x+5)}{(2x-1)(6x+3)}$		3	M1 factorise numerator M1 factorise denominator These marks are independent
		$\frac{x+5}{3(2x+1)}$		A1 $\frac{x+5}{6x+3}$
		3(2x+1)		Total 3 marks

17 (a)	$\frac{40}{32}$ or 75 or $\frac{5}{4}$ or $1\frac{1}{4}$ or 1.25		2	M1
		1 hour 15 minutes		A1
(b)	$\frac{1.5 + 40 + 10}{22.5 + 75 + 37.5} \times 60 = \frac{51.5}{135} \text{ or}$ $\frac{1.5 + 40 + 10}{0.375 + "1.25" + 0.625} = \frac{51.5}{2.25} \text{ (oe)}$		2	M1 FT 75 or 1.25 from (a)
		22.9		A1 awrt 22.9
				Total 4 marks

18	3x + y = 7	21x + 7y = 49	y = 7 - 3x	x =		4	M1	Two equations with equal	Allow a
	3x + 21y = -39	x + 7 v = -13		-13-7y				x or y coefficients, or	single
								correctly isolate <i>x</i> or <i>y</i> .	mistake
	20y = -46	20x = 62	<i>x</i> +	3 ×			M1	Correct choice to add or	across
			7(7-3x)	(-13-7y)			DEP	subtract equations, or	both M
			=-13	+ y = 7				substitute x or y into other	marks.
								equation	
					x = 3.1 (oe)		A1		
					y = -2.3 (oe)		A1		
								Tota	l 4 marks

19	$\sin 42 = \frac{5.2}{AC}$		4	M1	Any fully correct equation containing AC
	$AC = \frac{5.2}{\sin 42}$ (= 7.77 or better)			M1 DEP	Any fully correct expression for AC
	$\tan CAD = \frac{4.5}{"7.77"}$			M1 INDEP	ft $AC > 0$ Any fully correct expression for CAD or a trig ratio of CAD.
		30.1		A1	
					Total 4 marks
20 (a)		$2x - \frac{16}{x^2}$	2	B2	B1 for 2x, B1 for $-\frac{16}{x^2}$ or $-16x^{-2}$
(b)	$2x - \frac{16}{x^2} = 0$ (oe)		2	M1	$\frac{dy}{dx} = 0$, ft their $\frac{dy}{dx}$ as long as this is
		2		A1	not the same as y CSO
		_			Total 4 marks
21	6+2x=14		4	M1	
		x = 4		A1	
	$3y-3x = -27$ or $3y-3 \times "4" = -27$			M1	
		y = -5		A1	
	NB award both M marks if $\binom{6+2x}{3y-3x} = \binom{14}{-27}$	seen.			Total 4 marks

22	$Ext_{12} = \frac{360}{12}$ (=	30) $Int_{12} = \frac{(2 \times 12 - 4)90}{12}$		4	M1	
	12	(= 150)				
	$Ext_n = 54 - "30"$				M1	
		(= 156)			INDEP	
	$n = \frac{360}{"24"}$	$\frac{(2n-4)90}{n}$ = "156" (oe)			M1 DEP	Dependent on both previous method marks
		·	15		A1	
						Total 4 marks
23 (a)		135	1	B1	
(t		3)	133	3	M1	Allow different variable used.
	· ·	r-3=46 or (r-49)(r+3)=0 or	or		M1	
					(DEP)	
	(r=)	$\frac{(-46)^2 - 4 \times 1 \times -147}{2 \times 1}$				
			49		A1	49 and -3 together gain M1M1A0
	NB for factorise $46r - 147$ eg.	ation in (b) allow any expression w $(r-50)(r+4)$	which gives 2 out of 3 corr	ect in r^2 –		Total 4 marks
					,	
24 (a	$\begin{array}{c} 0.2 + x + x + 0. \\ 3x + 0.58 = 1 \end{array}$.1 + 0.18 + x + 0.1 = 1, be)		2	M1	
			0.14 (oe)		A1	$\frac{7}{50}$
(t)		2	1	B1	ft for $x > 0$
(0	e) 60×0.2			2	M1	
			12		A1	
						Total 5 marks

25	(a)	$2 \times 2.5 + 5 - (10)$			2	M1	Fully correct substitution into
		${2.5-3} - ({-0.5})$					function
				-20		A1	
	(b)	xy - 3y = 2x + 5	xy - 3x = 2y + 5		3	M1	
		x(y-2) = 3y + 5	y(x-2) = 3x + 5			M1	
						DEP	
				$\frac{3x+5}{x-2} \text{ (oe)}$		A1	$3 + \frac{11}{x - 2}$
							Total 5 marks

26	(a)	360 – 255 – (180 – 135) or 60 correct on diagram		2	M1	SC Answer of 300 M1A0
			60		A1	
						Accept 60(°), 060(°), N60(°)E
	(b)		5	1	B1	
	(c)	" 60 " – $(180-135)$ or $180-(360-255)$ – " 60 "		2	M1	
		(oe) or 15 correct on diagram				
			15		A1	Accept 15(°), 015(°), N15(°)E
	•					Total 5 marks

27	(a) (i)	equal intersecting arcs, centres A and C		4	M1	
			perpendicular bisector		A1	Any length.
			drawn			
	(ii)	arc centre A to intersect AC and AB , and equal			M1	
		intersecting arcs from these points of intersection				
			angle bisector drawn		A1	Must extend from <i>A</i> to <i>BC</i> .
	(b)	Arc, centre B, radius 5 cm		2	M1	
		A R B	region R shaded		A1 FT	FT their perpendicular bisector and angle bisector. The area must be bounded by at least both of these and the correct arc.
						Total 6 marks

28 (a) (i)	$2\pi r = \frac{216}{360} \times 2\pi \times 5x \text{ (oe)}$		4	M1
		3x		A1
(ii)	$(5x)^2 = ("3x")^2 + h^2 \text{ (oe)}$			M1
		4 <i>x</i>		A1
(b)	$\frac{1}{3}\pi("3x")^2 \times "4x" = \frac{375\pi}{2}$		3	M1 Condone missing brackets for this mark only
	$x^3 = \frac{375 \times 3}{2 \times 9 \times 4} \left(= \frac{125}{8} \right) $ (oe)			M1 DEP
		$2\frac{1}{2}$ (oe)		A1 $\frac{5}{2}$, 2.5
				Total 7 marks

