



Pearson

Mark Scheme (Results)

Summer 2017

Pearson Edexcel International GCSE
In Mathematics B (4MB0) Paper 02R

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2017

Publications Code 4MB0_02R_1706_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
 - cao – correct answer only
 - ft – follow through
 - isw – ignore subsequent working
 - SC - special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - awrt – answer which rounds to
 - eoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**


Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

| Question | Working | Answer | Mark | Notes |
|----------|---|---|------|---|
| 1 (a) | $\frac{360}{2} \times (2 + 3 + 5)$ | | | M1 |
| | | 1800 | 2 | A1 |
| (b) (i) | $(y =) \frac{3}{10} \times "1800"$ $(=540)$ | $(z =) \frac{5}{10} \times "1800"$ $(=900)$ | | M1 Calculation for either y or z Not retrospective (ie only award in (b) if used in (b)) |
| | "540" $\times 1.25$ | | | M1 (DEP) |
| | | 675 euros | | A1 |
| (ii) | "900" $\times 1.25 \times 1.2$ | | | M1 (DEP) |
| | (=1125 $\times 1.2$) | 1350 dollars | 5 | A1 |
| | | | | Total 7 marks |

| | | | | | |
|---|---|------|---|----------------------|--|
| 2 | $(2x+3)(x+1) = (3x-5)(x+2)$ | | | M1 | Any correct equation. |
| | $2x^2 + 3x + 2x + 3 = 3x^2 + 6x - 5x - 10$ | | | M1 (DEP) | Correctly expand either (quadratic) side. |
| | $x^2 - 4x - 13 (= 0)$ (oe) | | | A1 | cao |
| | $x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \times 1 \times (-13)}}{2 \times 1}$ (oe, completing the square ie “ $(x-2)^2 = 17$ ”) | | | M1 (INDEP) | ft if their quadratic has three non-zero terms. |
| | $x = \frac{4 \pm \sqrt{68}}{2}$ or $x = 2 \pm \sqrt{17}$ | | | M1 (DEP) | DEP on previous M1 ft for evaluating discriminant (ft if working seen and their discriminant is not negative.) |
| | NB: Some working must be seen else M0 M0 A0 if answer is incorrect. | 6.12 | 6 | A1 cao | DEP on third M1 Do not award if negative value is also given and not rejected. |
| | | | | Total 6 marks | |

| | | | | | |
|----------|---|--|----------------------------------|---|------------------------------------|
| 3 | (a) | | 90° | 1 | B1 cao |
| | (b) | | 45° | 1 | B1 cao |
| | (c) (i) | | 68° | | B1 cao |
| | | | corresponding angles | | B1 |
| | (ii) | | $\angle CEB = 45^\circ$ | | B1 cao |
| | | | angles in same segment | | B1 OR angles subtended by same arc |
| | (iii) | | $\angle EFB = 67^\circ$ | | B1 cao |
| | | | angles in triangle = 180° | 6 | B1 OR \angle s in Δ |
| | (d) | | 136° | 1 | B1 cao |
| | NB: (1) Award marks if angles seen on diagram (2) Reasons must be stated clearly and not to be inferred from their working. | | | | Total 9 marks |


| | | | | |
|----------|------------------------------|--|---|---|
| | Penalise nc ONCE only | | | |
| 4 | (a) | $AB^2 = 6^2 + 9^2 - 2 \times 6 \times 9 \cos 105$ | | M1 |
| | | $AB^2 = 117 - 108 \cos 105 (= 117 + 27.95 = 144.95)$ | | M1 (DEP) |
| | | 12.0 cm (Accept 12) | 3 | A1 12.03962... at least 3SF |
| | (b) | $\frac{1}{2} \times 6 \times 9 \sin 105$ | | M1 |
| | | 26.1 cm ² | 2 | A1 26.07999... at least 3 SF |
| | (c) | $\frac{352}{"26.1"}$ | | M1 |
| | | 13.5 | 2 | A1 13.49693... at least 3 SF |
| | (d) | $(6 + 9 + "12.0") \times "13.5" + 2 \times "26.1"$ | | M1 |
| | | 417 cm ² | 2 | A1 417.112... at least 3 SF (or awrt 417 if nc has been penalised) |
| | | | | Total 9 marks |

| | | | | |
|----------------------|---|--|---|--|
| 5 (a) | |  | 3 | <p>B3 B2 for 4 correct entries added B1 for 2 correct entries added</p> <p>NB: Start entering marks starting with the 1st B box so B2 (out of 3) would be entered as 1 1 0</p> |
| (b) | $x + 25 - x + 23 - x + 21 - x + 7 + x + 2 + x + x - 1 + 7 = 100$ (their 8 entries) | | | M1 ft Venn diagram |
| | | 16 (cao) | 2 | A1 |
| (c) | “ $23 - x + x + 21 - x$ ” or $(23 - 16) + 16 + (21 - 16)$ NB: Numbers can be on Venn Diagram | | | M1 ft Venn diagram Do not condone negative members numbers |
| | | 28 (cao) | 2 | A1 |
| (d) | | $\frac{5}{11}$ (cao) | 1 | B1 $\frac{25}{55}, 0.454\dots, 0.455, 45.4\%, 45.5\%$ |
| Total 8 marks | | | | |

| | | | | | | |
|---|-----|--|--------------------|---|---------------|---|
| 6 | (a) | $14 \times 24 + 17 \times 22 + 20 \times 28 + 24 \times 20 + 29 \times 6$ (= 1924) | | | M1 M1(DEP) | 3 correct fx products added Σfx for consistent x values in each interval (all correct) NB: Must use mid-values |
| | | $\frac{1924}{100}$ | | | M1 (DEP) | |
| | | | awrt 19.2 km/l | 4 | A1 | So accept, eg, 19.24 |
| | (b) | bars 16 – 18 height 11 cm (22 ss), 18 – 22 height 7 cm (14 ss), 26 – 32 height 1 cm (2 ss) | correct bars drawn | 3 | B3 | B1 for each bar with correct width and height. |
| | (c) | $\frac{1}{4} \times 20$ oe | | | M1 | Method to find the number of cars in the interval $25 < x \leq 26$ |
| | | | 11 (cao) | 2 | A1 | |
| | | NB: Thus 5 \rightarrow M1 A0 | | | | Total 9 marks |

| | | | | | | |
|---|-----|--|---|---|------------|--|
| 7 | (a) | | 2000 dollars | 1 | B1 | |
| | (b) | entrance fee = 1.2×8 (oe) (=9.6) visitors = 0.9×250 (oe) (=225) | | | M1 | Complete method to find new entrance fee OR number of visitors. |
| | | $1.2 \times 8 \times 0.9 \times 250$ | | | M1 (DEP) | |
| | | | 2160 dollars | 3 | A1 | |
| | (c) | entrance fee = $\frac{100+2r}{100} \times 8$ (=8 + 0.16r) | | | M1 (INDEP) | Complete method to find new entrance fee. |
| | | visitors = $\frac{100-r}{100} \times 250$ (=250-2.5r) | | | M1 (INDEP) | Complete method to find new number of visitors. |
| | | $(T \Rightarrow) \left(\frac{100+2r}{100} \times 8 \right) \left(\frac{100-r}{100} \times 250 \right)$ OR $(8 + 0.16r)(250 - 2.5r)$ OR $0.2(100+2r)(100-r)$ | | | M1 (DEP) | |
| | | $0.2(10000 - 100r + 200r - 2r^2)$ | $2000 + 20r - 0.4r^2$ (cso) | 4 | A1 | dep on M marks <u>Expansion of brackets must be shown.</u> |
| | (d) | $\frac{dT}{dr} = 20 - 0.8r$ | $-0.4(r^2 - 50r + 5000)$ (ie rewriting) | | M1 | One term correct |
| | | " $20 - 0.8r = 0$ " | $-0.4((r - 25)^2 + 5000 - 625)$ ie completing the square | | M1 (DEP) | Cand's derivative must be function of r |
| | | | 25 (cao) | 3 | A1 | |
| | | | | | | Total 11 marks |

| | | | | | |
|---|-----|--|--|---|--|
| 8 | (a) | | 3 0.75 | 2 | B1 B1 |
| | (b) | | Correct curve drawn | 3 | B3 –1 mark for each point missed/incorrectly plotted each point or segment missed straight line segments (penalise ONCE) tramlines (penalise ONCE) very poor curve NB: FT for (–1, ”3”) and (0.5, ”0.75”) Tol = $\pm \frac{1}{2}ss = \pm 0.025$ |
| | (c) | $2x^3 - 3x + 2 = 1\frac{1}{2}$ OR $2x^3 - 3x + 2 = -2x^3 + 3x + 1$ | | | M1 |
| | | | –1.3, 0.2, 1.1 (cao) Tol = $\pm 1ss = \pm 0.05$ | 3 | A2 A1 for two correct (So in ePEN this is scored at 1 then 0) |
| | (d) | $2x^3 - 3x + 2 > 1 - 4x$ | | | M1 condone “=” |
| | | Plot “ $y = 1 - 4x$ ” | | | M1(INDEP) ft (must be a straight line) |
| | | –0.6 | | | A1 ft from graph dep on above M1 |
| | | | $x > -0.6$ | 4 | A1 cao |
| | | | Tol = $\pm 1ss = \pm 0.05$ for both As above | | Total 12 marks |



| | | | | | |
|----------|---------|--|---|---|--|
| 9 | (a) (i) | | $4\mathbf{b}$ | | B1 |
| | (ii) | | $"4\mathbf{b}" + 8\mathbf{a}$ | | B1ft ft $4\mathbf{b}$ from (i) |
| | (iii) | | $4\mathbf{a} - 6\mathbf{b}$ | 3 | B1 |
| | (b) (i) | | $\lambda"(4\mathbf{a} - 6\mathbf{b})"$ | | B1ft ft $4\mathbf{a} - 6\mathbf{b}$ from (a)(iii) |
| | (ii) | $"4\mathbf{b}" + 4\mathbf{a} + "\lambda(4\mathbf{a} - 6\mathbf{b})"$ OR $-2\mathbf{b} + 8\mathbf{a} + (\lambda - 1)"(4\mathbf{a} - 6\mathbf{b})"$ | $4(\lambda + 1)\mathbf{a} + (4 - 6\lambda)\mathbf{b}$ | 2 | B1ft ft $\lambda(4\mathbf{a} - 6\mathbf{b})$ from (i) OR (a)(iii) Simplification NOT required. |
| | (c) | | $\mu"(4\mathbf{b}" + 8\mathbf{a})"$ | 1 | B1ft ft $4\mathbf{b} + 8\mathbf{a}$ from (a)(ii) |
| | (d) | $"4(\lambda + 1)\mathbf{a} + (4 - 6\lambda)\mathbf{b}" = "\mu(4\mathbf{b} + 8\mathbf{a})"$ $"4 + 4\lambda = 8\mu"$ and $"4 - 6\lambda = 4\mu"$ | | | M1 ft on (b)(ii) and (c) M1(DEP) ft |
| | | $-4 + 16\lambda = 0$ or $20 = 32\mu$ | | | M1 ft eliminate either variable |
| | | NB: If just one of $\lambda = \frac{1}{4}$ or $\mu = \frac{5}{8}$ and no working seen, score 4 / 5 marks | $\lambda = \frac{1}{4},$ $\mu = \frac{5}{8}$ | 5 | A1 cso A1 cso |
| | (e) (i) | | 20 | | B1 cao |
| | (ii) | | 5 | 2 | B1 cao |
| | | | | | Total 13 marks |

| | | | | | |
|---------------|---|--|---|----------------------|---|
| 10 (a) | $(2^{3x})^{x-2} = 2^{3x^2-6x}$ | | | M1 | Multiply indices |
| | $\frac{12 \times 6^{2x-1}}{9^x} = \frac{2^2 \times 3 \times 2^{2x-1} \times 3^{2x-1}}{(3^2)^x}$ <p>OR</p> $\frac{3 \times 2^2 \times 2^{3x(x-2)} \times 2^{2x-1} \times 3^{2x-1}}{3^{2x}}$ | | | M1 | Express 12, 6 and 9 as products of 2 and 3. |
| | | | | (M1) (M1(DEP)) | Factors of 2 OR 3 separated Factors of 2 AND 3 separated |
| | | $n = 3x^2 - 4x + 1$ Accept 2^{3x^2-4x+1} | 3 | A1 | cs0 |
| (b) | $3x^2 - 4x + 1 = 5$ | | | M1 | (really a B1) |
| | $(3x+2)(x-2) = 0$ | | | M1(INDEP) | or correct use of formula NB: Attempt on their 3 term quadratic |
| | | $x = -\frac{2}{3}$ and $x = 2$ (cao) | 3 | A1 | |
| | | | | Total 6 marks | |

| | | | | | | |
|-----------|-----|---|---|---|----------------|--|
| 11 | (a) | | (1, 2), (3, 2), (3, 1) | 1 | B1 | Coordinates or column vectors, any order |
| | (b) | | Rotation 270° centre (0, 0) | 3 | B1 B1 B1 | or -90° or 90° clockwise or <i>O</i> or origin |
| | (c) | | $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ | 1 | B1 | |
| | (d) | $\begin{pmatrix} -1 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 & 3 & 3 \\ 2 & 2 & 1 \end{pmatrix}$ NB: Order is important | | | M1 | fit from (a) for three correct entries Columns could be in any order. |
| | | | (-5, 4), (-7, 4), (-5, 2) | 2 | A1 cao | Accept $\begin{pmatrix} -5 & -7 & -5 \\ 4 & 4 & 2 \end{pmatrix}$ with columns in any order. |
| | (e) | $\begin{pmatrix} -1 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ NB: Order is important OR $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} -2 & -1 & -2 \\ 1 & 3 & 3 \end{pmatrix} = \begin{pmatrix} -5 & -5 & -7 \\ 4 & 2 & 4 \end{pmatrix}$ NB:(1) Order of coords important TWO correct eqns in <i>a</i> and <i>b</i> from “above” AND TWO correct eqns in <i>c</i> and <i>d</i> from “above” Answer (cao) | | | M1 | fit from (c) (M1) fit from (d) (M1 (DEP)) (A1) |

| | | | | |
|--|--|--|---|---|
| | | $\begin{pmatrix} 2 & -1 \\ -2 & 0 \end{pmatrix}$ | 3 | M1 ft their <i>correct</i> product of <i>their</i> 2x2 matrices A1 cao |
| | | | | Total 10 marks |

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom